

- Arneeth, A., et al. Comparative ecosystem–atmosphere exchange of energy and mass in a European Russian and a central Siberian bog II. Interseasonal and interannual variability of CO<sub>2</sub> fluxes. *Tellus B Chem. Phys. Meteorol.* **54**, 514–530 (2002).
- Aurela, M., et al. Carbon dioxide exchange on a northern boreal fen. *Boreal Environ. Res.* **14**, 699–710 (2009).
- Aurela, M., Laurila, T., & Tuovinen, J. P. The timing of snow melt controls the annual CO<sub>2</sub> balance in a subarctic fen. *Geophys. Res. Lett.* **31**, 3–6 (2004).
- Björkman, M. P. et. al. Temporal patterns of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O fluxes and soil microbial structure within snow-covered Alpine plant communities. (2007)
- Christiansen, C. T., Schmidt, N. M. & Michelsen, A. High Arctic dry heath CO<sub>2</sub> exchange during the early cold season. *Ecosystems* **15**, 1083-1092 (2012).
- Dolman, A. J., et al. Net ecosystem exchange of carbon dioxide and water of far eastern Siberian Larch (*Larix cajanderii*) on permafrost. *Biogeosciences* **1**, 133–146 (2004).
- Elberling, B. Annual soil CO<sub>2</sub> effluxes in the High Arctic: The role of snow thickness and vegetation type. *Soil Biol. Biochem.* **39** (2007).
- Elberling, B., Brandt, K.K. Uncoupling of microbial CO<sub>2</sub> production and release in frozen soil and its implications for field studies of arctic C cycling. *Soil Biol. Biogeochem.* **35**, 263-272 (2003).
- Euskirchen, E. S., Bret-Harte, M. S., Shaver, G. R., Edgar, C. W., Romanovsky, V. E. . Long-term release of carbon dioxide from arctic tundra ecosystems in Alaska. *Ecosystems* **20**, 960–974 (2017).

- Euskirchen, E. S., Bret-Harte, M. S., Scott, G. J., Edgar, C., & Shaver, G. R. Seasonal patterns of carbon dioxide and water fluxes in three representative tundra ecosystems in northern Alaska. *Ecosphere* **3** (2012).
- Fahnestock, J. T., Jones, M. H., Brooks, P. D., Walker, D. A., & Welker, J. M. Winter and early spring CO<sub>2</sub> efflux from tundra communities of northern Alaska. *J. Geophys. Res. Atmos.* **103**, 29023–29027 (1998).
- Grogan, P. Cold season respiration across a Low Arctic landscape: The influence of vegetation type, snow depth, and interannual climatic variation. *Arctic, Antarct. Alp. Res.* **44**, 446–456 (2012).
- Grogan, P., & Jonasson, S. Ecosystem CO<sub>2</sub> production during winter in a Swedish subarctic region: the relative importance of climate and vegetation type. *Glob. Chang. Biol.* **12**, 1479–1495 (2006).
- Grogan, P., & Jonasson, S. Temperature and substrate controls on intra-annual variation in ecosystem respiration in two subarctic vegetation types. *Glob. Chang. Biol.* **11**, 465–475 (2005).
- Grogan, P., & Chapin, F. S. Arctic soil respiration: Effects of climate and vegetation depend on season. *Ecosystems* **2**, 451–459 (1999).
- Heikkinen, J. E. P., Elsakov, V., & Martikainen, P. J. Carbon dioxide and methane dynamics and annual carbon balance in tundra wetland in NE Europe, Russia. *Global Biogeochem. Cycles* **16** (2002).
- Helbig, M., et al. Direct and indirect climate change effects on carbon dioxide fluxes in a thawing boreal forest–wetland landscape. *Glob. Chang. Biol.* **23**, 3231–3248 (2017).

- Ives, S. L., Sullivan, P. F., Dial, R., Berg, E. E., & Welker, J. M. CO<sub>2</sub> exchange along a hydrologic gradient in the Kenai Lowlands, AK: Feedback implications of wetland drying and vegetation succession. *6*, 38–50 (2013).
- Joiner, D. W., Lafleur, P. M., McCaughey, J. H., & Bartlett, P. A. Interannual variability in carbon dioxide exchanges at a boreal wetland in the BOREAS northern study area. *J. Geophys. Res. Biogeosciences* **104**, 27663–27672 (1999).
- Jones, M. H., Fahnesstock, J. T., & Welker, J. M. Early and late winter CO<sub>2</sub> efflux from Arctic tundra in the Kuparuk River watershed, Alaska. *Arctic, Antarct. Alp. Res.* **31**, 187–190 (1999).
- Kade, A., Bret-Harte, M. S., Euskirchen, E. S. E. S., Edgar, C., & Fulweber, R. A. . Upscaling of CO<sub>2</sub> fluxes from heterogeneous tundra plant communities in Arctic Alaska. *J. Geophys. Res. Biogeosciences* **117**, 1-11 (2012).
- Kim, Y., Y. Kodama, and G.J. Fochesatto. Environmental factors regulating winter CO<sub>2</sub> flux in snow-covered black forest soil of Interior Alaska. *Geochemical Journal*, (51), 2017.
- Kim, Y., Park, S. J., Lee, B. Y., & Risk, D. Continuous measurement of soil carbon efflux with Forced Diffusion (FD) chambers in a tundra ecosystem of Alaska. *Sci. Total Environ.* **566**, 175–184 (2016).
- Kim, Y. Effect of ablation rings and soil temperature on 3-year spring CO<sub>2</sub> efflux along the Dalton Highway, Alaska. *Biogeosciences* **11**, 6539–6552 (2014).
- Kim, Y., et al. Latitudinal distribution of soil CO<sub>2</sub> efflux and temperature along the Dalton Highway, Alaska. *Polar Sci.* **7**, 162–173 (2013).
- Kim, Y., et al. Assessment of winter fluxes of CO<sub>2</sub> and CH<sub>4</sub> in boreal forest soils of central Alaska estimated by the profile method and the chamber method: A diagnosis of methane

- emission and implications for the regional carbon budget. *Tellus, Ser. B Chem. Phys. Meteorol.* **59**, 223–233 (2007).
- Larsen, K. S., Grogan, P., Jonasson, S., & Michelsen, A. Respiration and microbial dynamics in two subarctic ecosystems during winter and spring thaw: Effects of increased snow depth. *Arctic, Antarct. Alp. Res.* **39**, 268-276 (2007).
- Larsen, K. S., Ibrom, A., Grogan, P., Jonasson, S., Michelsen, A., & Beier, C. Significance of cold-season respiration and photosynthesis in a subarctic heath ecosystem in Northern Sweden. *Global Change Biology*, 2007.
- Luërs, J., et al. Annual CO<sub>2</sub> budget and seasonal CO<sub>2</sub> exchange signals at a high Arctic permafrost site on Spitsbergen, Svalbard archipelago. *Biogeosciences* **11**, 6307–6322 (2014).
- Morgner, E., Elberling, B., Strebel, D., & Cooper, E. J. The importance of winter in annual ecosystem respiration in the High Arctic: Effects of snow depth in two vegetation types. *Polar Res.* **29**, 58–74 (2010).
- Natali, S. M., Schuur, E.A.G., Trucco, C., Hicks Pries, C.E., et al. Effects of experimental warming of air, soil and permafrost on carbon balance in Alaskan tundra. *Glob. Chan. Biol.* **17**, 1394-1407 (2011).
- Natali, S. M., Schuur, E. A. G. G., Webb, E. E., Pries, C. E. H. H., & Crummer, K. G. Permafrost degradation stimulates carbon loss from experimentally warmed tundra. *Ecology* **95**, 602–608 (2014).
- Nobrega, S., & Grogan, P. Deeper snow enhances winter respiration from both plant-associated and bulk soil carbon pools in birch hummock tundra. *Ecosystems* **10**, 419–431 (2007).

- Nordstroem, C., Soegaard, H., Christensen, T. R., Friborg, T., & Hansen, B. U. Seasonal carbon dioxide balance and respiration of a high-arctic fen ecosystem in NE-Greenland. *Theor. Appl. Climatol.* **70**, 149–166 (2001).
- Oechel, W. C., et al. A scaling approach for quantifying the net CO<sub>2</sub> flux of the Kuparuk River Basin, Alaska. *Glob. Chang. Biol.* **6**, 160–173 (2000).
- Oechel, W. C., et al. Acclimation of ecosystem CO<sub>2</sub> exchange in the Alaskan Arctic in response to decadal climate warming. *Nature* **406**, 978–981 (2000).
- Olefeldt, D., et al. Net carbon accumulation of a high-latitude permafrost palsa mire similar to permafrost-free peatlands. *Geophys. Res. Lett.* **39** (2012).
- Pirk, N., et al. Snowpack fluxes of methane and carbon dioxide from high Arctic tundra. *J. Geophys. Res. Biogeosciences* **121**, 2886–2900 (2016).
- Rogers, M. C., Sullivan, P.F., Welker, J.M. Evidence of nonlinearity in the response of net ecosystem CO<sub>2</sub> exchange to increasing levels of winter snow depth in the high Arctic of Northwestern Greenland. *Arct. Antarc. Alpine Res.* **43**, 95-106 (2011).
- Schimel, J. P., Bilbrough, C., & Welker, J. M. Increased snow depth affects microbial activity and nitrogen mineralization in two Arctic tundra communities. *Soil Biol. Biochem.* **36**, 217–227 (2004).
- Semenchuk, P. R., Christiansen, C. T., Grogan, P., Elberling, B., & Cooper, E. J. Long-term experimentally deepened snow decreases growing-season respiration in a low- and high-arctic tundra ecosystem. *J. Geophys. Res. Biogeosciences* **121**, 1236–1248 (2016).

- Sjögersten, S., van der Wal, R., & Woodin, S. J. Habitat type determines herbivory controls over CO<sub>2</sub> fluxes in a warmer Arctic. *Ecology* **89**, 2103–2116 (2008).
- Strebel, D., Elberling, B., Morgner, E., Knicker, H. E., & Cooper, E. J. Cold-season soil respiration in response to grazing and warming in High-Arctic Svalbard. *Polar Res.* **29**, 46–57 (2010).
- Sullivan, P. F., Welker, J. M., Arens, S. J. T. T., Sveinbjornsson, B., & Sveinbjörnsson, B. Continuous estimates of CO<sub>2</sub> efflux from arctic and boreal soils during the snow-covered season in Alaska. *J. Geophys. Res. Biogeosciences* **113** (2008).
- Sullivan, P. F. Snow distribution, soil temperature and late winter CO<sub>2</sub> efflux from soils near the Arctic treeline in northwest Alaska. *Biogeochemistry* **99**, 65–77 (2010).
- Trucco, C., et al. Seven-year trends of CO<sub>2</sub> exchange in a tundra ecosystem affected by long-term permafrost thaw. *J. Geophys. Res. Biogeosciences* **117** (2012).
- Ueyama, M., Iwata, H., & Harazono, Y. Autumn warming reduces the CO<sub>2</sub> sink of a black spruce forest in interior Alaska based on a nine-year eddy covariance measurement. . *Glob. Chang. Biol.* **20**, 1161–1173 (2014).
- Vogel, J., Schuur, E. A. G., Trucco, C., & Lee, H. Response of CO<sub>2</sub> exchange in a tussock tundra ecosystem to permafrost thaw and thermokarst development. *J. Geophys. Res. Biogeosciences* **114** (2009).
- Vogel, J. G., Valentine, D. W., & Ruess, R. W. Soil and root respiration in mature Alaskan black spruce forests that vary in soil organic matter decomposition rates. *Can. J. For. Res.* **35**, 161–174 (2005).

- Webb, E. E., et. al. Increased wintertime CO<sub>2</sub> loss as a result of sustained tundra warming. *Biogeosciences* **121**, 1-17 (2016).
- Welker, J. M., Fahnestock, J. T., Henry, G. H. R., O'Dea, K. W., & Chimner, R. A. CO<sub>2</sub> exchange in three Canadian High Arctic ecosystems: Response to long-term experimental warming. *Glob. Chang. Biol.* **10**, 1981–1995 (2004).
- Welker, J. M., Fahnestock, J.T., Jones, M.H. Annual CO<sub>2</sub> flux in dry and moist arctic tundra: field responses to increases in summer temperatures and winter snow depth. *Clim. Chan* **44**, 139-150 (2000).
- Welp, L. R., Randerson, J. T., & Liu, H. P. The sensitivity of carbon fluxes to spring warming and summer drought depends on plant functional type in boreal forest ecosystems. *Agric. For. Meteorol.* **147**, 72–185 (2007).
- Wickland, K. P., Striegl, R. G., Neff, J. C., & Sachs, T. Effects of permafrost melting on CO<sub>2</sub> and CH<sub>4</sub> exchange of a poorly drained black spruce lowland. *J. Geophys. Res.* *Biogeosciences* **111**, 1–13 (2006).
- Winston, G. C., Sundquist, E. T., & Stephens, B. B. Winter CO<sub>2</sub> fluxes in a boreal forest. *J. Geophys. Res. Biogeosciences* **102**, 795–804 (1997).
- Zimov, S. A. Winter biotic activity and production of CO<sub>2</sub> in Siberian soils: a factor in the greenhouse effect. *J. Geophys. Res. Atmos.* **98**, 5017–5023 (1993).
- Zimov, S. A., et al. Siberian CO<sub>2</sub> efflux in winter as a CO<sub>2</sub> source and cause of seasonality in atmospheric CO<sub>2</sub>. *Clim. Change* **33**, 111–120 (1996).