

References to the Root Profiles Data Set

References followed by a (*) mark were used in the study by Schenk, H. J. and R. B. Jackson.
2002. The global biogeography of roots. Ecological Monographs. 72(3):311-328.

- Abaimov, A. P., S. G. Prokushkin, O. A. Zyryanova, and L. N. Kaverzina. 1997. Osobennosti formirovaniya i funkcionirovaniya listvenichnykh lesov na merzlotnykh pochvakh. (Peculiarities of forming and functioning larch forests on frozen soils. In Russian). Lesovedenie 1997(5):13-23. (*)
- Aerts, R. 1993. Biomass and nutrient dynamics of dominant plant species from heathlands. Pages 51-84 in R. Aerts, and G. W. Heil, editors. Heathlands: patterns and processes in a changing environment. Kluwer, Dordrecht, The Netherlands. (*)
- Alvarez-Lopez, M. 1990. Ecology of *Pterocarpus officinalis* forested wetlands in Puerto Rico. Pages 251-265 in A. E. Lugo, M. Brinson, and S. Brown, editors. Forested wetlands. Elsevier, Amsterdam.
- Ares, A., and N. Peinemann. 1992. Fine-root distribution of coniferous plantations in relation to site in southern Buenos Aires, Argentina. Canadian Journal of Forest Research 22:1575-1582. (*)
- Bang-xing, W. 1991. Studies on the vertical structure of seasonal rain-forest in Xishuangbanna of Yunnan. Acta Botanica Sinica 33:232-239. (*)
- Barbour, M. G., J. A. MacMahon, S. A. Bamberg, and J. A. Ludwig. 1977. The structure and distribution of *Larrea* communities. Pages 227-251 in T. J. Mabry, J. H. Hunziker, and D. R. DiFeo, Jr., editors. Creosote bush: biology and chemistry of *Larrea* in New World deserts. US/IBP Series 6. Dowden, Hutchinson and Ross, Stroudsburg, Pennsylvania, USA. (*)
- Bard, G. E. 1952. Secondary succession on the Piedmont of New Jersey. Ecological Monographs 22:195-215.
- Barker, D. J., D. M. Zhang, and A. D. Mackay. 1988. Root distribution in a low fertility hill country sward grazed by sheep. New Zealand Journal of Experimental Agriculture 16:73-76. (*)
- Belsky, A. J. 1994. Influences of trees on savanna productivity: tests of shade, nutrients, and tree-grass competition. Ecology 75:932. (*)
- Berish, C. W. 1982. Root biomass and surface area in three successional tropical forests. Canadian Journal of Forest Research 12:699-704. (*)
- Berish, C. W., and J. J. Ewel. 1988. Root development in simple and complex tropical successional ecosystems. Plant and Soil 106:73-84. (*)

- Bernard, J. M., and K. Fiala. 1986. Distribution and standing crop of living and dead roots in three wetland Carex species. *Bulletin of the Torrey Botanical Club* 113:1-5.
- Bhatti, J. S., N. W. Foster, and P. W. Hazlett. 1998. Fine root biomass and nutrient content in a black spruce peat soil with and without alder. *Canadian Journal of Soil Science* 78:163-169. (*)
- Bille, J. C. 1977. Étude de la production primaire nette d'un écosystème Sahélien. *Travaux et Documents de l'O.R.S.T.O.M.* 65. O.R.S.T.O.M., Paris, France. (*)
- Blagoveshchenskiy, E. N. 1968. The dry savanna of northwest India. *Soviet Geography* 9:519-537. (*)
- Boikov, T. G., and Y. D. Kharitonov. 1998. Biomass of underground organs in Transbaikalian steppe phytocenoses. Pages 55-60 in J. E. Box, Jr., editor. *Root demographics and their efficiencies in sustainable agriculture, grasslands and forest ecosystems. Developments in Plant and Soil Sciences* 82. Kluwer, Dordrecht, The Netherlands. (*)
- Bowden, B. N. 1963. The root distribution of *Andropogon gayanus* var. *bisquamulatus*. *East African Agricultural and Forestry Journal* 29:157-159.
- Bowns, J. E., and N. E. West. 1976. Blackbrush (*Coleogyne ramosissima* Torr.) on southwestern Utah rangelands. *Research Report* 27. Utah Agricultural Experiment Station, Logan, Utah, USA. (*)
- Bowns, J. E., Jr., and T. W. Box. 1964. The influence of grazing on the roots and rhizomes of seacoast bluestem. *Journal of Range Management* 17:36-39.
- Branson, F. A., R. F. Miller, and I. S. McQueen. 1976. Moisture relationships in twelve northern desert shrub communities near Grand Junction, Colorado. *Ecology* 57:1104-1124. (*)
- Briones, O., C. Montaña, and E. Ezcurra. 1996. Competition between three Chihuahuan desert species: evidence from plant size-distance relations and root distribution. *Journal of Vegetation Science* 7:453-460. (*)
- Carbon, B. A., G. A. Bartle, A. M. Murray, and D. K. MacPherson. 1980. The distribution of root length, and the limits to flow of soil water to roots in a dry sclerophyll forest. *Forest Science* 26:656-664. (*)
- Castellanos, J., M. Maass, and J. Kummerow. 1991. Root biomass of a dry deciduous tropical forest in Mexico. *Plant and Soil* 131:225-228. (*)
- Cavelier, J. 1992. Fine-root biomass and soil properties in a semideciduous and a lower montane rain forest in Panama. *Plant and Soil* 142:187-201. (*)

- Cerri, C. C., and B. Volkoff. 1987. Carbon content in a yellow Latosol of central Amazon rain forest. *Acta Oecologica. Oecologia Generalis* 8:29-42. (*)
- Chapman, S. B. 1970. The nutrient content of the soil and root system of a dry heath ecosystem. *Journal of Ecology* 58:445-452. (*)
- Chen, Z., H. Chang, and B. Wang. 1994. Studies on biomass and production of the lower subtropical evergreen broad-leaved forest in Heishiding nature reserve, China. VI. Distribution, biomass and production of roots. *Journal of Tropical Ecology* 10:273-279. (*)
- Christie, E. K. 1978. Ecosystem processes in semiarid grasslands. I Primary production and water use of two communities possessing different photosynthetic pathways. *Australian Journal of Agricultural Research* 29:773-787. (*)
- Cline, J. F., D. W. Uresk, and W. H. Rickard. 1977. Plants and soil of a sagebrush community on the Hanford Reservation. *Northwest Science* 51:60-70.
- Coetzee, J. A., M. I. Page, and D. Meredith. 1946. Root studies in Highveld grassland communities. *South African Journal of Science* 42:105-118.
- Coughenour, M. B., J. E. Ellis, and R. G. Popp. 1990. Morphometric relationships and developmental patterns of *Acacia tortilis* and *Acacia reficiens* in Southern Turkana, Kenya. *Bulletin of the Torrey Botanical Club* 117:8-17. (*)
- Coupland, R. T., and T. C. Brayshaw. 1953. The fescue grassland in Saskatchewan. *Ecology* 34:386-405. (*)
- Coupland, R. T., J. R. Willard, E. A. Ripley, and R. L. Randell. 1975. The Matador Project. Pages 19-50 in T. W. M. Cameron, and L. W. Billingsley, editors. *Energy flow - its biological dimensions*. Royal Society of Canada, Ottawa, Canada. (*)
- Cox, J. R., G. W. Frasier, and K. G. Renard. 1986. Biomass distribution at grassland and shrubland sites. *Rangelands* 8:67-69.
- Daddy, F., M. J. Trlica, and C. D. Bonham. 1988. Vegetation and soil water differences among big sagebrush communities with different grazing histories. *Southwestern Naturalist* 33:413-424. (*)
- Dahlman, R. C., and C. L. Kucera. 1965. Root productivity and turnover in native prairie. *Ecology* 46:84-89. (*)
- Damman, A. W. H. 1971. Effect of vegetation changes on the fertility of a Newfoundland forest site. *Ecological Monographs* 41:253-270. (*)
- Davies, S. J., and P. Becker. 1996. Floristic composition and stand structure of mixed dipterocarp and heath forests in Brunei Darussalam. *Journal of Tropical Forest Science* 8:542-569. (*)

- Davis, E. A., and C. P. Pase. 1977. Root system of shrub live oak: Implications for water yield in Arizona chaparral. *Journal of Soil and Water Conservation* 32:174-180. (*)
- Davis, G. R., W. A. Neilsen, and J. G. McDavis. 1983. Root distribution of *Pinus radiata* related to soil characteristics in five Tasmanian soils. *Australian Journal of Soil Research* 21:165-171. (*)
- de Castro, E. A., and J. B. Kauffman. 1998. Ecosystem structure in the Brazilian Cerrado: a vegetation gradient of aboveground biomass, root mass and consumption by fire. *Journal of Tropical Ecology* 14:263-283. (*)
- Dennis, J. G., and P. L. Johnson. 1970. Shoot and rhizome-root standing crops of tundra vegetation at Barrow, Alaska. *Arctic and Alpine Research* 2:253-266. (*)
- Dennis, J. G., L. L. Tieszen, and M. A. Vetter. 1978. Seasonal dynamics of above- and belowground production of vascular plants at Barrow, Alaska. Pages 113-140 in L. L. Tieszen, editor. *Vegetation and production ecology of an Alaskan arctic tundra*. Ecological Studies 29. Springer-Verlag, New York, New York, USA. (*)
- Devidas, S., and J.-P. Puyravaud. 1995. Primary productivity of the herbaceous layer in a grazed savanna woodland, Bandipur National Park, southern India. *Acta Oecologica* 16:491-505. (*)
- Dhyani, S. K., P. Narain, and R. K. Singh. 1990. Studies on root distributions of five multipurpose tree species in Doon Valley, India. *Agroforestry Systems* 12:149-161.
- Distel, R. A., and O. A. Fernández. 1988. Dynamics of root growth and decay in two grasses native to semi-arid Argentina. *Australian Journal of Ecology* 13:327-336. (*)
- Dobrowolski, J. P., M. M. Caldwell, and J. H. Richards. 1990. Basin hydrology and plant root systems. Pages 243-292 in C. B. Osmond, L. F. Pitelka, and G. M. Hidy, editors. *Plant biology of the basin and range*. Ecological Studies 80. Springer-Verlag, Berlin, Germany. (*)
- Dumortier, M. 1991. Below-ground dynamics in a wet grassland ecosystem. Pages 301-309 in D. Atkinson, editor. *Plant root growth: an ecological perspective*. Blackwell Scientific Publications, Oxford.
- Duncan, W. H. 1941. A study of root development in three soil types in the Duke Forest. *Ecological Monographs* 11:141-164. (*)
- Ellenberg, H., R. Mayer, and J. Schauermann, editors. 1986. *Ökosystemforschung. Ergebnisse des Sollingprojekts 1966-1986*. Ulmer, Stuttgart, Germany. (*)
- Evdokimova, T. I., and L. A. Grishina. 1968. Productivity of root systems of herbaceous vegetation on flood plain meadows and methods for its study. Pages 24-27 in M. S. Ghilarov, V. A. Kovda, L. N. Novichkova-Ivanova, L. E. Rodin, and V. M. Sveshnikova, editors. *Methods of*

- productivity studies in root systems and rhizosphere organisms. International Symposium USSR, August 28--September 12, 1968. Nauka, Leningrad, Russia. (*)
- Farrish, K. W. 1991. Spatial and temporal fine-root distribution in three Louisiana forest soils. *Soil Science Society of America Journal* 55:1752-1757. (*)
- Fernandez, O. A., and M. M. Caldwell. 1975. Phenology and dynamics of root growth of three cool semi-desert shrubs under field conditions. *Journal of Ecology* 63:703-714. (*)
- Fernández, R. J., and J. M. Paruelo. 1988. Root systems of two Patagonian shrubs: A quantitative description using a geometrical method. *Journal of Range Management* 41:220-223. (*)
- Fiala, K. 1990. Live and dead underground plant biomass in a natural meadow hydrosere. *Folia Geobotanica and Phytotaxonomica* 25:113-135. (*)
- Fiala, K., and R. Herrera. 1988. Living and dead belowground biomass and its distribution in some savanna communities in Cuba. *Folia Geobotanica and Phytotaxonomica* 23:225-237. (*)
- Fiala, K., and V. Studený. 1988. Cutting and fertilization effect on the root system in several grassland stands. II. Vertical distribution of root biomass and changes in the carbohydrate content. *Ekológia* 7:27-42. (*)
- Finér, L., C. Messier, and L. De Grandpré. 1997. Fine-root dynamics in mixed boreal conifer--broad-leaved forest stands at different successional stages after fire. *Canadian Journal of Forest Research* 27:304-314. (*)
- Frangi, J. L., and A. E. Lugo. 1985. Ecosystem dynamics of a subtropical floodplain forest. *Ecological Monographs* 55:351-369.
- Freckman, D. W., and R. A. Virginia. 1989. Plant-feeding nematodes in deep-rooting desert ecosystems. *Ecology* 70:1665-1678. (*)
- Friesen, D. K., I. M. Rao, R. J. Thomas, A. Oberson, and J. I. Sanz. 1997. Phosphorus acquisition and cycling in crop and pasture in low fertility tropical soils. *Plant and Soil* 196:289-294. (*)
- Garelkov, D. 1973. Biological productivity of some beech forest types in Bulgaria. Pages 307-314 in H. E. Young, editor. IUFRO biomass studies. College of Sciences and Agriculture, University of Maine, Orono, Maine, USA. (*)
- Gaze, S. R., J. Brouwer, L. P. Simmonds, and J. Bromley. 1998. Dry season water use patterns under *Guiera senegalensis* L. shrubs in a tropical savanna. *Journal of Arid Environments* 40:53-67. (*)
- Gehrmann, J., M. Gerriets, J. Puhe, and B. Ulrich. 1984. Untersuchungen an Boden, Wurzeln, Nadeln und erste Ergebnisse von Depositionsmessungen im Hils. Berichte des Forschungszentrums Waldökosysteme 2. (*)

- Gisi, U., and J. J. Oertli. 1981. Oekologische Entwicklung in Brachland verglichen mit Kulturwiesen. *Acta Oecologica. Oecologia Plantarum* 2:79-86. (*)
- Glatzel, G. 1983. Root distribution and soil water depletion in an oak-hornbeam stand (*Quercus petraea*, *Q. robur*, *Carpinus betulus*) and a spruce thicket (*Picea abies*). Pages 577-584 in W. Böhm, L. Kutschera, and E. Lichtenegger, editors. *Wurzelökologie und ihre Nutzanwendung/Root ecology and its practical application*. Bundesanstalt für alpenländische Landwirtschaft Grumpenstein, Irdning, F. R. Germany. (*)
- Gower, S. T. 1987. Relations between mineral nutrient availability and fine root biomass in two Costa Rica wet forests: a hypothesis. *Biotropica* 19:171-175. (*)
- Greenland, D. J., and J. M. L. Kowal. 1960. Nutrient content of the moist tropical forest of Ghana. *Plant and Soil* 12:154-174. (*)
- Greenwood, K. L., and K. J. Hutchinson. 1998. Root characteristics of temperate pasture in New South Wales after grazing at three stocking rates for 30 years. *Grass and Forage Science* 53:120-128. (*)
- Groeneveld, D. P. 1989. Shrub rooting and water acquisition on threatened shallow groundwater habitats in the Owens Valley, California. Pages 221-237 in *Proceedings--Symposium on cheatgrass invasion, shrub die-off, and other aspects of shrub biology and management*. General Technical Report INT-276. U.S.D.A. Forest Service Intermountain Research Station, Ogden, Utah, USA. (*)
- Groot, J. J. R., D. Koné, M. Traoré, and N. Kamissoko. 1995. Description du système racinaire de l'*Andropogon gayanus*, du *Vigna unguiculata* et du *Stylosanthes hamata* en zone soudano-sahélienne. *Rapports PSS 8. Production Soudano-Sahélienne (PSS)*, Wageningen, The Netherlands. (*)
- Groves, R. H., and R. L. Specht. 1965. Growth of heath vegetation I. Annual growth curves of two heath ecosystems in Australia. *Australian Journal of Botany* 13:261-280. (*)
- Hansson, A.-C., A. Zhao, and O. Andrén. 1995. Fine-root production and mortality in degraded vegetation in Horqin sand rangeland in Inner Mongolia, China. *Arid Soil Research and Rehabilitation* 9:1-13. (*)
- Harris, W. F., R. S. Kinerson, Jr., and N. T. Edwards. 1977. Comparison of belowground biomass of natural deciduous forest and loblolly pine plantations. *Pedobiologia* 17:369-381. (*)
- Haworth, F. 1953. Observations on the root system of Guatemala grass on an upland tea soil in Ceylon. *Tropical Agriculture* 30:116-121.

Heitschmidt, R. K., R. J. Ansley, S. L. Dowhower, P. W. Jacoby, and D. L. Price. 1988. Some observations from the excavation of honey mesquite root systems. *Journal of Range Management* 41:227-231. (*)

Hendriks, C. M. A., and F. J. J. A. Bianchi. 1995. Root density and root biomass in pure and mixed forest stands of Douglas-fir and beech. *Netherlands Journal of Agricultural Science* 43:321-331. (*)

Hertel, D. 1999. Das Feinwurzelsystem von Rein- und Mischbeständen der Rotbuche: Struktur, Dynamik und interspezifische Konkurrenz. *Dissertationes Botanicae* 317. J. Cramer, Berlin, Germany. (*)

Higgins, K. B., A. J. Lamb, and B. W. van Wilgen. 1987. Root systems of selected plant species in mesic mountain fynbos in the Jonkershoek Valley, south-western Cape Province. *South African Journal of Botany* 53:249-257. (*)

Hosegood, P. H. 1963. The root distribution of Kikuyu grass and Wattle trees. *East African Agricultural and Forestry Journal* 29:60-61. (*)

Hulbert, L. C. 1955. Ecological studies of *Bromus tectorum* and other annual bromegrasses. *Ecological Monographs* 25:181-213. (*)

Huttel, C. 1975. Root distribution and biomass in three Ivory Coast rain forest plots. Pages 123-130 in F. B. Golley, and E. Medina, editors. *Tropical ecological systems: Trends in terrestrial and aquatic research*. Ecological Studies 11. Springer-Verlag, Heidelberg, Germany. (*)

Ignatenko, I. V., A. V. Knorre, N. V. Lovelius, and B. N. Norin. 1972. Standing crop in plant communities at the station Ary-Mas. Pages 140-148 in F. E. Wielgolaski, and T. Rosswall, editors. *Tundra biome. International Biological Programme (IBP) Tundra Biome Steering Committee*, Stockholm, Sweden. (*)

Ignatenko, I. V., and F. I. Khakimzyanova. 1971. Soils and total phytomass reserves in dwarf birch - white dryas and willow tundras of the east European northlands. *Soviet Journal of Ecology* 2:300-305. (*)

Isagi, Y., T. Kawahara, K. Kamo, and H. Ito. 1997. Net production and carbon cycling in a bamboo *Phyllostachys pubescens* stand. *Plant Ecology* 130:41-52. (*)

Jackson, R. B., J. L. Banner, E. Jobbág, W. T. Pockman, and D. H. Wall. 2002. Ecosystem carbon loss with woody plant invasion of grasslands. *Nature* 418:623-626.

Jama, B., R. J. Buresh, J. K. Ndifa, and K. D. Shepherd. 1998. Vertical distribution of roots and soil nitrate: tree species and phosphorus effects. *Soil Science Society of America Journal* 62:280-286. (*)

Joffre, R., M. J. Leiva Morales, S. Rambal, and R. Fernández Ales. 1987. Dynamique racinaire et extraction de l'eau du sol par des graminées pérennes et annuelles méditerranéennes. *Acta Oecologica Oecologia Plantarum* 8:181-194. (*)

Johnsen, T. N., Jr. 1962. One-seed juniper invasion of northern Arizona grasslands. *Ecological Monographs* 2:187-207. (*)

Jonsson, K., L. Fidjeland, J. A. Maghembe, and P. Höglberg. 1988. The vertical distribution of roots of five tree species and maize in Mogoro, Tanzania. *Agroforestry Systems* 6:63-69.

Jordan, P. W., and P. S. Nobel. 1984. Thermal and water relations of roots of desert succulents. *Annals of Botany* 54:705-718. (*)

Kaletkina, N. G. 1974. Sezonnoe razvitiye rastitel'nosti subal'pijskoj raznotravnoj stepi i kriofil'ioj pustoshi Gissarskogo chrebita. (Seasonal development of vegetation in the subalpine steppe and cryophytic heath of the Hissar Mountains. In Russian). Pages 7-41 in P. N. Ovchinnikov, editor. *Rastitel'nost' Tadzhikistana i ee osvoenie*. (Vegetation of Tajikistan and its use). Donish, Dushanbe. (*)

Kalisz, P. J., R. W. Zimmerman, and R. N. Muller. 1987. Root density, abundance, and distribution in the mixed mesophytic forest of eastern Kentucky. *Soil Science Society of America Journal* 51:220-225. (*)

Kandeler, E., G. Eder, and M. Sobotik. 1994. Microbial biomass, N mineralization, and the activities of various enzymes in relation to nitrate leaching and root distribution in a slurry-amended grassland. *Biology and Fertility of Soils* 18:7-12.

Karizumi, N. 1978. Underground biomass. Pages 82-88 in T. Kira, Y. Ono, and T. Hosokawa, editors. *Biological production in a warm-temperate evergreen oak forest of Japan*. JIBP Synthesis 18. University of Tokyo Press, Tokyo, Japan. (*)

Karpov, V. G., editor 1983. *Faktory reguliatsii ekosistem elovykh lesov* (Regulation factors of spruce forest ecosystems. In Russian). Nauka, Leningrad, Russia. (*)

Karpov, V. G., editor. 1983. *Faktory reguliatsii ekosistem elovykh lesov* (Regulation factors of spruce forest ecosystems. In Russian). Nauka, Leningrad.

Kellman, M., and K. Sanmugadas. 1985. Nutrient retention by savanna ecosystems I. Retention in the absence of fire. *Journal of Ecology* 73:935-951. (*)

Kellman, M., and N. Roulet. 1990. Nutrient flux and retention in a tropical sand-dune succession. *Journal of Ecology* 78:664-676. (*)

Kern, K. G., W. Moll, and H. J. Braun. 1961. Wurzeluntersuchungen in Rein- und Mischbeständen des Hochschwarzwaldes (Vfl. Todtmoos 2/I-IV). (The rooting in pure and mixed stands of the

upper Black Forest, experimental plot Todtmoos 2/I-IV. In German with English summary). Allgemeine Forst- und Jagdzeitung 132:241-260. (*)

Khodachek, E. A. 1971. Vegetal matter of tundra phytocoenoses in the western part of Taymyr Peninsula (Translated from the original Russian publication in Botanicheskii Zhurnal 54:1059-1073). International Tundra Biome Translation 5. Tundra Biome Center, University of Alaska, College, Alaska, USA. (*)

Kimmins, J. P., and B. C. Hawkes. 1978. Distribution and chemistry of fine roots in a white spruce - subalpine fir stand in British Columbia: implications for management. Canadian Journal of Forest Research 8:265-279. (*)

Klepper, E. L., K. A. Gano, and L. L. Cadwell. 1985. Rooting depth and distributions of deep-rooted plants in the 200 Area Conrol Zone of the Hanford site. Pacific Northwest Laboratory, Richland, Washington.

Klinge, H. 1973. Root mass estimation in lowland tropical rain forests of central Amazonia, Brazil. I. Fine root masses of pale yellow Latosol and a giant humus Podzol. Tropical Ecology 14:29-38. (*)

Klinge, H. 1975. Root mass estimation in lowland tropical rain forests of central Amazonia, Brazil. III. Nutrients in fine roots from giant humus Podsols. Tropical Ecology 16:28-38. (*)

Knoop, W. T., and B. H. Walker. 1985. Interactions of woody and herbaceous vegetation in a Southern African savanna. Journal of Ecology 73:235-253. (*)

Kochenderfer, J. N. 1973. Root distribution under some forest types native to West Virginia. Ecology 54:445-448. (*)

Kosmas, C. S., N. Moustakas, N. G. Danalatos, and N. Yassoglou. 1996. The Spata field site: I. The impacts of land use and management on soil properties and erosion. II. The effect of reduced soil moisture on soil properties and wheat production. Pages 207-228 in C. J. Brandt, and J. B. Thornes, editors. Mediterranean desertification and land use. John Wiley and Sons, Chichester, UK. (*)

Kreutzer, K. 1968. The root system of the red alder (*Alnus glutinosa* Gärtn.). Pages 114-119 in M. S. Ghilarov, V. A. Kovda, L. N. Novichkova-Ivanova, L. E. Rodin, and V. M. Sveshnikova, editors. Methods of productivity studies in root systems and rhizosphere organisms. International Symposium USSR, August 28--September 12, 1968. Nauka, Leningrad, Russia. (*)

Kudrjasheva, O. I. 1974. Sezonnoe razvitiie rastitel'nosti nizkotravnnykh polusavann v juzhnoj okonechnosti chrebeta Aruk-Tau (stacionar Garauty). (Seasonal development of vegetation in the shortgrass steppe at the southern end of the Aruk-Tau Mountains (Station Garavuti). In Russian). Pages 77-105 in P. N. Ovchinnikov, editor. Rastitel'nost' Tadzhikistana i ee osvoenie. (Vegetation of Tajikistan and its use). Donish, Dushanbe, Tajikistan. (*)

Kul'tiasov, M. V. 1925. Materialy po izucheniju isparenija i kornevoj sistemy soobshchestva vesehhich zfemerov. (Materials on a study of evaporation and the root system of an association of spring ephemerals. In Russian with French summary). Biulleten' Sredne-Aziatskogo Gosudarstvennogo Universiteta 10:79-87. (*)

Kummerow, J., and R. Mangan. 1981. Root systems in *Quercus dumosa* Nutt. dominated chaparral in southern California. Acta Oecologica. Oecologia Plantarum 2:177-188. (*)

Kummerow, J., M. Kummerow, and L. Trabaud. 1990. Root biomass, root distribution and the fine-root growth dynamics of *Quercus coccifera* L. in the garrigue of southern France. Vegetatio 87:37-44. (*)

Ladygina, G. M., and N. P. Litvinova. 1974. Produktivnost' nekotorych lugovych soobshchestv vostochnogo Pamira. Problemy Botaniki 12:275-285. (*)

Lamont, B. 1973. Factors affecting the distribution of protiod roots within the root systems of two *Hakea* species. Australian Journal of Botany 21:165-187. (*)

Lavrinenko, D. D. 1972. Interaction of wood species in different types of forests. Indian National Scientific Documentation Center, New Delhi, India. (*)

Lawson, G. W., J. Jenik, and K. O. Armstrong-Mensah. 1968. A study of a vegetation catena in Guinea savanna at Mole Game Reserve (Ghana). Journal of Ecology 56:505-522. (*)

Lawson, G. W., K. O. Armstrong-Mensah, and J. B. Hall. 1970. A catena in tropical moist semi-deciduous forest near Kade, Ghana. Journal of Ecology 58:371-398. (*)

Le Roux, X., T. Bariac, and A. Mariotti. 1995. Spatial partitioning of the soil water resource between grass and shrub components in a West African humid savanna. Oecologia 104:147-155. (*)

Lee, C. A., and W. K. Lauenroth. 1994. Spatial distribution of grass and shrub root systems in the shortgrass steppe. American Midland Naturalist 132:117-123. (*)

Liang, Y. M., D. L. Hazlett, and W. K. Lauenroth. 1989. Biomass dynamics and water use efficiencies of five plant communities in the shortgrass steppe. Oecologia 80:148-153. (*)

Linkola, K., and A. Tiirkka. 1936. Über Wurzelsysteme und Wurzelabspreitung der Wiesenpflanzen auf verschiedenen Wiesenstandorten. Annales Botanici Societatis Zoologicae Botanicae-Fennicae Vanamo Vol. 6, No. 6. Druckerei der Finnischen Literatur- Gesellschaft, Helsinki, Finland. (*)

Low, A. B., and B. B. Lamont. 1990. Aerial and below-ground phytomass of Banksia scrub-heath at Eneabba, South-western Australia. Australian Journal of Botany 38:351-359. (*)

- Lucot, E., and S. Bruckert. 1992. Organisation du système racinaire du chêne pédonculé (*Quercus robur*) développé en conditions édaphiques non contraignantes (sol brun lessivé colluvial). Annales des Sciences Forestières 49:465-479. (*)
- Lutz, H. J. L., J. B. Ely, Jr., and S. L. Little, Jr. 1937. The influence of soil profile horizons on root distribution of white pine (*Pinus strobus* L.). Yale University School of Forestry Bulletin 44:1-75. (*)
- Manning, S. J., and M. G. Barbour. 1988. Root systems, spatial patterns, and competition for soil moisture between two desert shrubs. American Journal of Botany 75:885-893. (*)
- Martínez, F., O. Merino, A. Martín, D. García Martín, and J. Merino. 1998. Belowground structure and production in a Mediterranean sand dune shrub community. Plant and Soil 201:209-216. (*)
- McClaugherthy, C. A., J. D. Aber, and J. M. Melillo. 1982. The role of fine roots in the organic matter and nitrogen budgets of two forested ecosystems. Ecology 63:1481-1490. (*)
- McKell, C. M., M. B. Jones, and E. R. Perrier. 1962. Root production and accumulation of root material on fertilized annual range. Agronomy Journal 54:459-462. (*)
- McNaughton, S. J., F. F. Banyikwa, and M. M. McNaughton. 1998. Root biomass and productivity in a grazing ecosystem: the Serengeti. Ecology 79:587-592. (*)
- Mekkonen, K., R. J. Buresh, and B. Jama. 1997. Root and inorganic nitrogen distributions in sesbania fallow, natural fallow and maize fields. Plant and Soil 188:319-327. (*)
- Mensah, K. O. A., and J. Jenik. 1968. Root system of tropical tress 2. Features of the root system of iroko (*Chlorophora excelsa* Benth. et Hook.). Preslia 40:21-27. (*)
- Metsävainio, K. 1931. Untersuchungen über das Wurzelsystem der Moorplanten. Druckerei der Finnischen Literatur- Gesellschaft, Helsinki.
- Midwood, A. J., T. W. Boutton, S. R. Archer, and S. E. Watts. 1998. Water use by woody plants on contrasting soils in a savanna parkland: assessment with d₂H and d₁₈O. Plant and Soil 205:13-24. (*)
- Miller, P. C., and E. Ng. 1977. Root:shoot biomass ratios in shrubs in southern California and central Chile. Madroño 24:215-223. (*)
- Miller, P. C., R. Mangan, and J. Kummerow. 1982. Vertical distribution of organic matter in eight vegetation types near Eagle Summit, Alaska. Holarctic Ecology 5:117-124. (*)
- Millikin, C. S., and C. S. Bledsoe. 1999. Biomass and distribution of fine and coarse roots from blue oak (*Quercus douglasii*) trees in the northern Sierra Nevada foothills of California. Plant and Soil 214:27-38.

- Miroshnichenko, Y. M. 1975. Kornevye sistemy drevesnych i kustarnikovych rastenij i ich zkologija v vostocnych karakumach. (Root systems of trees and bushes and their ecology in eastern Karakum. In Russian). Botanicheskii Zhurnal 60:1776-1795. (*)
- Montaña, C., B. Cavagnaro, and O. Briones. 1995. Soil water use by co-existing shrubs and grasses in the southern Chihuahuan Desert, Mexico. Journal of Arid Environments 31:1-13. (*)
- Moorhead, D. L., J. F. Reynolds, and P. J. Fonteyn. 1989. Patterns of stratified soil water loss in a Chihuahuan Desert community. Soil Science 148:244-249. (*)
- Mordelet, P., J.-C. Menaut, and A. Mariotti. 1997. Tree and grass rooting patterns in an African humid savanna. Journal of Vegetation Science 8:65-70. (*)
- Moreira, M. Z., L. d. S. L. Sternberg, and D. Nepstad. 2000. Vertical patterns of soil water uptake by plants in a primary forest and an abandoned pasture in the eastern Amazon: an isotopic approach. Plant and Soil 222:95-107.
- Muc, M. 1977. Ecology and primary production of sedge-moss meadow communities, Truelove Lowland. Pages 157-184 in L. C. Bliss, editor. Truelove Lowland, Devon Island, Canada: A high arctic ecosystem. University of Alberta Press, Edmonton, Alberta, Canada. (*)
- Murphy, P. G., A. E. Lugo, A. J. Murphy, and D. C. Nepstad. 1995. The dry forests of Puerto Rico's south coast. Pages 178-209 in A. E. Lugo, and C. Lowe, editors. Tropical forests: management and ecology. Ecological Studies 112. Springer-Verlag, New York, New York, USA. (*)
- Nepstad, D. C., C. R. de Carvalho, E. A. Davidson, P. H. Jipp, P. A. Lefebvre, G. H. Negreiros, E. D. da Silva, T. A. Stone, S. E. Trumbore, and S. Vieira. 1994. The role of deep roots in the hydrological and carbon cycles of Amazonian forests and pastures. Nature 372:666-669. (*)
- Nnyamah, J. U., and T. A. Black. 1977. Rates and patterns of water uptake in a Douglas-fir forest. Soil Science Society of America Journal 41:972-979. (*)
- Nobel, P. S. 1989. Temperature, water availability, and nutrient levels at various soil depths: Consequences for shallow-rooted desert succulents, including nurse plant effects. American Journal of Botany 76:1486-1492. (*)
- Nobel, P. S., M. E. Loik, and R. W. Meyer. 1991. Microhabitat and diel tissue acidity changes for two sympatric cactus species differing in growth habit. Journal of Ecology 79:167-182. (*)
- Odum, H. T. 1970. Rain forest structure and mineral-cycling homeostasis. Pages H-3--H-52 in H. T. Odum, and R. F. Pigeon, editors. A tropical rain forest: A study of irradiation and ecology at El Verde, Puerto Rico. Division of Technical Information, U.S. Atomic Energy Commission, Oak Ridge, Tennessee, USA. (*)

- Okali, D. U. U., J. B. Hall, and G. W. Lawson. 1973. Root distribution under a thicket clump on the Accra Plains, Ghana: Its relevance to clump localization and water relations. *Journal of Ecology* 61:439-454. (*)
- Old, S. M. 1969. Microclimate, fire, and plant production in an Illinois prairie. *Ecological Monographs* 39:355-384. (*)
- Olsthoorn, A. F. M. 1991. Fine root density and root biomass of two Douglas-fir stands on sandy soils in the Netherlands. 1. Root biomass in early summer. *Netherlands Journal of Agricultural Science* 39:49-60. (*)
- Pandey, C. B., and J. S. Singh. 1992. Influence of rainfall and grazing on belowground biomass dynamics in a dry tropical savanna. *Canadian Journal of Botany* 70:1885-1890. (*)
- Parker, M. M., and D. H. Van Lear. 1996. Soil heterogeneity and root distribution of mature loblolly pine stands in Piedmont soils. *Soil Science Society of America Journal* 60:1920-1925. (*)
- Pavlis, J., and Jeník.J. 2000. Roots of pioneer trees in the Amazonian rain forest. *Trees* 14:442-455.
- Pavón, N. P., and O. Briones. 2000. Root distribution, standing crop biomass and belowground productivity in a semidesert in Mexico. *Plant Ecology* 146:131-136.
- Persson, H. 1982. Changes in the tree and dwarf shrub fine-roots after clear-cutting in a mature Scots pine stand. *Swedish Coniferous Forest Project Technical Report* 31:19pp. (*)
- Persson, H., Y. von Fircks, H. Majdi, and L. O. Nilsson. 1995. Root distribution in a Norway spruce (*Picea abies* (L.) Karst.) stand subjected to drought and ammonium-sulfate application. *Plant and Soil* 168-169: 161-165. (*)
- Pietikäinen, J., E. Vaijärvi, H. Ilvesniemi, H. Fritze, and C. J. Westman. 1999. Carbon storage of microbes and roots and the flux of CO₂ across a moisture gradient. *Canadian Journal of Forest Research* 29:1197-1203. (*)
- Plamboeck, A. H., H. Grip, and U. Nygren. 1999. A hydrological tracer study of water uptake depth in a Scots pine forest under two different water regimes. *Oecologia* 119:452-460. (*)
- Plewczynska-Kuras, U. 976. Estimation of biomass of the underground parts of meadow herbage in the three variants of fertilization. *Polish Ecological Studies* 2:63-74. (*)
- Popov, K. P. 1979. Fistashka v srednei Azii. (The pistachio in central Asia. In Russian). Ylym, Ashkhabad, Turkmenistan. (*)
- Puhe, J., H. Persson, and I. Börjesson. 1986. Wurzelwachstum und Wurzelschäden in skandinavischen Nadelwäldern. (Root growth and root damages in Scandinavian coniferous forests. In German with English summary). *Allgemeine Forstzeitung* 20:488-492. (*)

- Qiu, G.-K., Y.-K. Shen, D.-Y. Li, Z.-W. Wang, Q.-M. Huang, D.-D. Yang, and A.-X. Gao. 1992. Bamboo in sub-tropical eastern China. Pages 159-188 in S. P. Long, M. B. Jones, and M. J. Roberts, editors. Primary productivity of grass ecosystems of the tropics and sub-tropics. Chapman and Hall, London, UK. (*)
- Qiu, X., S. Xie, and G. Jin. 1984. A preliminary study on biomass of *Lithocarpus xylocarpus* forest in Xujiaba region, Ailao mountains, Yunnan (China). *Acta Botanica Yunnanica* 6:85-92. (*)
- Raich, J. W. 1983. Effects of forest conversion on the carbon budget of a tropical soil. *Biotropica* 15:177-184. (*)
- Ramam, S. S. 1970. Root development in alluvial grasslands of Varanasi. *Indian Forester* 96:100-110.
- Rao, I. M. 1998. Root distribution and production in native and introduced pastures in the South American savannas. Pages 19-41 in J. E. Box, Jr., editor. Root demographics and their efficiencies in sustainable agriculture, grasslands and forest ecosystems. Kluwer Academic Publishers, Dordrecht, The Netherlands. (*)
- Reynolds, E. R. C. 1970. Root distribution and the cause of its spatial variability in *Pseudotsuga taxifolia* (Poir.) Britt. *Plant and Soil* 32:501-517. (*)
- Rickard, W. H., and B. E. Vaughan. 1988. Plant community characteristics and responses. Pages 109-179 in W. H. Rickard, L. E. Rogers, B. E. Vaughan, and S. F. Liebetrau, editors. Shrub-steppe: Balance and change in a semi-arid terrestrial ecosystem. Developments in Agricultural and Managed-Forest Ecology 20. Elsevier, Amsterdam, The Netherlands. (*)
- Roberts, J. 1976. A study of root distribution and growth in a *Pinus sylvestris* L. (Scots pine) plantation in East Anglia. *Plant and Soil* 44:607-621. (*)
- Rodin, L. E., editor 1977. *Produktivnost' rastitel'nosti aridnoi zony Azii (itogi sovetskikh issledovanij po Mezhdunarodnoj biologicheskoj programme, 1965-1977 gg.).* (Productivity of vegetation in arid zone of the Asia. (Synthesis of the Soviet studies for the International Biological Programme, 1965-1974. In Russian). Nauka, Leningrad, Russia. (*)
- Rosswall, T., J. G. K. Flower-Ellis, L. G. Johansson, S. Jonsson, B. E. Rydén, and M. Sonesson. 1975. Stordalen (Abisko), Sweden. Pages 265-294 in T. Rosswall, and O. W. Heal, editors. Structure and function of tundra ecosystems. Ecological Bulletin 20. Swedish Natural Science Research Council, Stockholm, Sweden. (*)
- Roupsard, O., A. Ferhi, A. Granier, F. Pallo, D. Depommier, B. Mallet, H. I. Joly, and E. Dreyer. 1999. Reverse phenology and dry-season water uptake by *Faidherbia albida* (Del.) A. Chev. in an agroforestry parkland of Sudanese west Africa. *Functional Ecology* 13:460-472.

- Rui, L., M. J. A. Werger, H. J. During, and Z. C. Zhong. 1999. Biomass distribution in a grove of the giant bamboo *Phyllostachys pubescens* in Chongqing, China. *Flora (Jena)* 194:89-96. (*)
- Saarinen, T. 1996. Biomass and production of two vascular plants in a boreal mesotrophic fen. *Canadian Journal of Botany* 74:934-938.
- Safford, L. O. 1974. Effect of fertilization on biomass and nutrient content of fine roots in a beech-birch-maple stand. *Plant and Soil* 40:349-363. (*)
- Safford, L. O., and S. Bell. 1972. Biomass of fine roots in a white spruce plantation. *Canadian Journal of Forest Research* 2:169-172. (*)
- Samoilova, E. M. 1968. The study of the tree root systems on sandy soils. Pages 195-200 in M. S. Ghilarov, V. A. Kovda, L. N. Novichkova-Ivanova, L. E. Rodin, and V. M. Sveshnikova, editors. *Methods of productivity studies in root systems and rhizosphere organisms. International Symposium USSR, August 28--September 12, 1968.* Nauka, Leningrad, Russia. (*)
- Sanford, R. L., Jr. 1989. Root systems of three adjacent, old growth Amazon forests and associated transition zones. *Journal of Tropical Forest Science* 1:268-279. (*)
- Saurina, N. I., and I. V. Kamenetskaya. 1969. Massa kornej sosny obyknovennoj (*Pinus sylvestris* L.) v sosnjake mshisto-lishajnikovom juzhnog tajgi. (The root mass of *Pinus sylvestris* L. in moss-lichen pine forests of the southern taiga. In Russian with English summary). *Bulleten Moskovskogo obshchestva ispytatelei prirody. Otdel biologicheskii* 74:96-100. (*)
- Scherfose, V. 1990. Feinwurzelverteilung und Mykorrhizatypen von *Pinus sylvestris* in verschiedenen Bodentypen. *Berichte des Forschungszentrums Waldökosysteme A* 62:166pp.
- Scholes, R. J., and B. H. Walker. 1993. *An African savanna: synthesis of the Nyala study. Cambridge Studies in Applied Ecology and Resource Management* Cambridge University Press, Cambridge, UK. (*)
- Schreiber, K.-F., A. Yair, and M. Shachak. 1995. Ecological gradients along slopes of the northern Negev highlands, Israel. Pages 209-229 in H.-P. Blume, and S. M. Berkowicz, editors. *Arid ecosystems. Advances in GeoEcology* 28. Catena Verlag, Cremlingen, Germany. (*)
- Schroth, G., D. Kolbe, B. Pity, and W. Zech. 1996. Root system characteristics with agroforestry relevance of nine leguminous tree species and a spontaneous fallow in a semi-deciduous rainforest area of West Africa. *Forest Ecology and Management* 84:199-208. (*)
- Schulze, E.-D., H. A. Mooney, O. E. Sala, E. Jobbagy, N. Buchmann, G. Bauer, J. Canadell, R. B. Jackson, J. Loreti, M. Oesterheld, and J. R. Ehleringer. 1996. Rooting depth, water availability, and vegetation cover along an aridity gradient in Patagonia. *Oecologia* 108:503-511. (*)
- Scully, N. J. 1942. Root distribution and environment in a maple-oak forest. *Botanical Gazette* 103:492-517. (*)

Shackleton, C. M., B. McKenzie, and J. E. Granger. 1988. Seasonal changes in root biomass, root/shoot ratios and turnover in two coastal grassland communities in Transkei. South African Journal of Botany 54:465-471. (*)

Shalyt, M. S. 1950. Podzemnaja cast' nekotorykh lugovykh, stepnykh i pustynnykh rasteniy i fitocenozov. C. I. Travjanistye i polukustarnikovye rastenija i fitocenozy lesnoj (luga) i stepnoj zon. (Belowground parts of some meadow, steppe, and desert plants and plant communities. Part I: Herbaceous plants and subshrubs and plant communities of forest and steppe zones. In Russian). Trudy Botanicheskogo Instituta im. V.L. Komarova. Akademii nauk SSSR. Seriia III, Geobotanika 6:205-442. (*)

Shalyt, M. S. 1952. Podzemnaja cast' nekotorykh lugovykh, stepnykh i pustynnykh rasteniy i fitocenozov. C. 2. Travjanistye, polukustarnikovye i kustarnikovye rastenija i fitocenozy pustynnoj zony. (Belowground parts of some meadow, steppe, and desert plants and plant communities. Part 2: Herbaceous plants, subshrubs, and shrubs, and plant communities of the desert zone. In Russian). Trudy Botanicheskogo Instituta im. V.L. Komarova. Akademii nauk SSSR. Seriia III, Geobotanika 8:71-139. (*)

Shalyt, M. S., and L. F. Zhivotenko. 1968. Overground and underground parts of certain grass and dwarf semishrub phytocoenoses of the Crimean Jaila (mountain pastures) and the technique of their estimation. Pages 204-208 in M. S. Ghilarov, V. A. Kovda, L. N. Novichkova-Ivanova, L. E. Rodin, and V. M. Sveshnikova, editors. Methods of productivity studies in root systems and rhizosphere organisms. International Symposium USSR, August 28--September 12, 1968. Nauka, Leningrad, Russia. (*)

Sims, P. L., J. S. Singh, and W. K. Lauenroth. 1978. The structure and function of ten western North American grasslands: II. Intra-seasonal dynamics in primary producer compartments. Journal of Ecology 66:547-572. (*)

Singh, J. S., and D. C. Coleman. 1977. Evaluation of functional root biomass and translocation of photoassimilated C14 in a shortgrass prairie ecosystem. Pages 29-37 in J. K. Marshall, editor. The belowground ecosystem: a synthesis of plant-associated processes. Range Science Department Science Series 26. Colorado State University, Fort Collins, Colorado, USA. (*)

Singh, V. 1994. Morphology and pattern of root distribution in *Prosopis cineraria*, *Dalbergia sissoo* and *Albizia lebbek* in an arid region of north-western India. Tropical Ecology 35:133-146. (*)

Sjörs, H. 1991. Phyto- and necromass above and below ground in a fen. Holarctic Ecology 14:208-218.

Smit, G. N., and N. F. G. Rethman. 1998. Root biomass, depth distribution and relations with leaf biomass of *Colophospermum mopane*. South African Journal of Botany 64:38-43. (*)

Sommer, R., M. Denich, and P. L. G. Vlek. 2000. Carbon storage and root penetration in deep soils under small-farmer land-use systems in the Eastern Amazon region, Brazil. *Plant and Soil* 210:231-241.

Soumaré, A., J. J. R. Groot, D. Koné, and S. Radersma. 1994. Structure spatiale du système racinaire de deux arbres du Sahel: *Acacia seyal* et *Sclerocarya birrea*. Rapports PSS 5. Production Soudano-Sahélienne (PSS), Wageningen, The Netherlands. (*)

Specht, R. L. 1957. Dark Island heath (Ninety-mile Plain, South Australia) III. The root system. *Australian Journal of Botany* 5:103-114. (*)

Srivastava, S. K., K. P. Singh, and R. S. Upadhyay. 1986. Fine root growth dynamics in teak (*Tectona grandis* Linn. F.). *Canadian Journal of Forest Research* 16:1360-1364. (*)

Steele, S. J., S. T. Gower, J. G. Vogel, and J. M. Norman. 1997. Root mass, net primary production and turnover in aspen, jack pine and black spruce forests in Saskatchewan and Manitoba, Canada. *Tree Physiology* 17:577-587. (*)

Sternberg, P. D., M. A. Anderson, R. C. Graham, J. L. Beyers, and K. R. Tice. 1996. Root distribution and seasonal water status in weathered granitic bedrock under chaparral. *Geoderma* 72:89-98. (*)

Strong, W. L., and G. H. La Roi. 1983. Rooting depths and successional development of selected boreal forest communities. *Canadian Journal of Forest Research* 13:577-588. (*)

Sturges, D. L. 1977. Soil water withdrawal and root characteristics of big sagebrush. *American Midland Naturalist* 98:257-273.

Sturges, D. L. 1980. Soil water withdrawal and root distribution under grubbed, sprayed, and undisturbed big sagebrush vegetation. *Great Basin Naturalist* 40:157-164. (*)

Sveshnikova, V. M. 1968. Root system distribution and biomass of plants in Pamir Mountain deserts. Pages 208-213 in M. S. Ghilarov, V. A. Kovda, L. N. Novichkova-Ivanova, L. E. Rodin, and V. M. Sveshnikova, editors. *Methods of productivity studies in root systems and rhizosphere organisms*. International Symposium USSR, August 28--September 12, 1968. Nauka, Leningrad, Russia. (*)

Symbula, M., and F. P. Day, Jr. 1988. Evaluation of two methods for estimating belowground production in a freshwater swamp forest. *American Midland Naturalist* 120:405-415.

Thompson, J., J. Proctor, V. Viana, W. Miliken, J. A. Ratter, and D. A. Scott. 1992. Ecological studies on a lowland evergreen rain forest on Maracá Island, Roraima, Brazil. I. Physical environment, forest structure and leaf chemistry. *Journal of Ecology* 80:689-703. (*)

Toky, O. P., and R. P. Bisht. 1992. Observations on the rooting patterns of some agroforestry trees in an arid region of north-western India. *Agroforestry Systems* 18:245-263. (*)

- Tryon, P. R., and F. S. Chapin, III. 1983. Temperature control over root growth and root biomass in taiga forest trees. Canadian Journal of Forest Research 13:827-833. (*)
- Turner, L. M. 1936. A comparison of roots of southern shortleaf pine in three soils. Ecology 17:649-658. (*)
- Uchida, M., T. Nakatsubo, T. Horikoshi, and K. Nakane. 1998. Contribution of micro-organisms to the carbon dynamics in black spruce (*Picea mariana*) forest soil in Canada. Ecological Research 13:17-26. (*)
- Usol'tsev, V. A., and I. S. Krepkii. 1994. Regression analysis of vertical-fraction distribution of root mass in Aman-Karagai pine forests. Russian Journal of Ecology 25:87-97. (*)
- Van Rees, K. C. J., and N. B. Comerford. 1986. Vertical root distribution and strontium uptake of a slash pine stand on a Florida spodosol. Soil Science Society of America Journal 50:1042-1046. (*)
- Vance, E. D., and N. M. Nadkarni. 1992. Root biomass distribution in a moist tropical montane forest. Plant and Soil 142:31-39. (*)
- Vandenbeldt, R. J. 1991. Rooting systems of western and southern African *Faidherbia albida* (Del.) A. Chev. (syn. *Acacia albida* Del.) - a comparative analysis with biogeographic implications. Agroforestry Systems 14:233-244. (*)
- Villares, J. B., A. Tundisi, and M. Becker. 1953. The subterranean system of colonial grass (Guinea grass) in various soils of the State of São Paulo, Brazil. Journal of Range Management 6:248-254.
- Vyskot, M. 1973. Root biomass of silver fir (*Abies alba* Mill.). Acta Universitatis Agriculturae (Brno), Series C 42:215-261. (*)
- Wallace, A., E. M. Romney, and J. W. Cha. 1980. Depth distribution of roots of some perennial plants in the Nevada Test Site area of the northern Mojave Desert. Great Basin Naturalist Memoirs 4:201-207. (*)
- Watts, S. E. 1993. Rooting patterns of co-occurring woody plants on contrasting soils in a subtropical savanna. Master's thesis. Texas A&M University, College Station, Texas, USA. (*)
- Weaver, J. E., and R. W. Darland. 1949. Soil-root relationships of certain native grasses in various soil types. Ecological Monographs 19:303-338. (*)
- Weaver, J. E., V. H. Hougen, and M. D. Weldon. 1935. Relation of root distribution to organic matter in prairie soil. Botanical Gazette 96:389-420. (*)

- Weaver, T. 1977. Root distribution and soil water regimes in nine habitat types of the northern Rocky Mountains. Pages 239-244 in J. K. Marshall, editor. The belowground ecosystem: a synthesis of plant-associated processes. Range Science Department Science Series 26. Colorado State University, Fort Collins, Colorado, USA. (*)
- Webber, P. J., and D. E. May. 1977. The magnitude and distribution of belowground plant structures in the alpine tundra of Niwot Ridge, Colorado. Arctic and Alpine Research 9:157-174. (*)
- Wright, T. W. 1955. Profile development in the sand dunes of Culbin Forest, Morayshire. Journal of Soil Science 6:270-283. (*)
- Xu, Y. 1991. Ökologische Grundlagen für den Anbau der Großen Küstentanne (*Abies grandis* Lindl.) auf vernässten Böden. Berichte des Forschungszentrums Waldökosysteme A 67: (*)
- Yano, N., and R. Kayama. 1975. Seasonal and yearly change of biomass and litter -- underground. Pages 147-160 in M. Numata, editor. Ecological studies in Japanese grasslands. JIBP Synthesis 13. University of Tokyo Press, Tokyo, Japan. (*)
- Yeager, A. F. 1935. Root systems of certain trees and shrubs grown on prairie soil. Journal of Agricultural Research 51:1085-1092.
- Yin, X., J. A. Perry, and R. K. Dixon. 1989. Fine-root dynamics and biomass distribution in a *Quercus* ecosystem following harvesting. Forest Ecology and Management 27:159-177. (*)
- Zhang Kebin. 1989. The growth of man-made forests of *Haloxylon ammodendron* and their soil water contents in the Minqin desert region, Gansu Province, China. Journal of Arid Environments 17:109-115. (*)
- Zverev, N. E., and R. D. Seiidova. 1990. Underground mass of shrub and subshrub plants of the Karakum. Problems of Desert Development 1990(1):49-54. (*)