


Search ORNL DAAC



 Search

[DAAC Home](#) >
 [Get Data](#) >
 [NASA Projects](#) >
 [MODIS/ASTER Airborne Simulator \(MASTER\)](#) >
 User guide

MASTER: Western Diversity Time Series Campaign, WDTS, California, USA, Spring 2021

Get Data

Documentation Revision Date: 2023-04-11

Dataset Version: 1.1


Summary

This dataset includes Level 1B (L1B) and Level 2 (L2) data products from the MODIS/ASTER Airborne Simulator (MASTER) instrument. The spectral data were collected as part of the Western Diversity Time Series (WDTS, formerly HypSIIRI) airborne campaign during nine flights aboard a NASA ER-2 aircraft over selected areas of California and Nevada, U.S., from 2021-02-09 to 2021-04-02. The WDTS campaign will observe California's ecosystems and provide critical information on natural disasters such as volcanoes, wildfires, and drought. MASTER products can identify vegetation type and health and provide a benchmark for the state of the ecosystems against which future changes can be assessed. Data products include L1B georeferenced multispectral imagery of calibrated radiance in 50 bands covering wavelengths of 0.460 to 12.879 micrometers at approximately 50-meter spatial resolution. Derived L2 data products are emissivity in 5 bands in thermal infrared range (8.58 to 12.13 micrometers) and land surface temperature. The L1B file format is HDF-4, and L2 products are provided in ENVI and KMZ formats. In addition, the dataset includes the flight path, spectral band information, instrument configuration, ancillary notes, and summary information for each flight, and browse images derived from each L1B data file.

The MASTER instrument is a modified Daedalus Wildfire scanning spectrometer that flies on a variety of multi-altitude research aircraft and provides spectral information similar to that provided by the Moderate Resolution Imaging Spectroradiometer (MODIS) and the Advanced Spaceborne Thermal Emission and Reflection Radiometer (ASTER), which are aboard two NASA Earth Observing System satellites: Terra and Aqua. The primary goal of this airborne campaign was to demonstrate important science and applications research that is uniquely enabled by the full suite of MASTER thermal infrared bands as well as the contiguous spectroscopic measurements of the AVIRIS (also flown in similar campaigns), or combinations of measurements from both instruments.

This dataset includes a total of 680 data files: 82 files in Hierarchical Data Format (HDF-4; *.hdf) format, 308 ENVI raster files (*.dat and *.hdr) that are compressed (*.zip), 77 files in Keyhole Markup Language Zipped (KMZ; *.kmz) format, 77 Portable Network Graphics (PNG; *.png) files that are compressed (*.zip), 36 text (*.txt) files, 9 archives of text files that are zipped (*.zip), 9 flight maps as GIF (*.gif) images, and 82 browse images in JPEG (*.jpg) format.

ASTER/MODIS Airborne Simulator Browse Imagery
 WDTS_Spring21 Campaign - 24 Feb 2021
 Ivanpah Playa (CA/NV)
 Flight #21-930-00 Track #3



Upper Left Lat, Lon = 35.8°, -115.6°
 Lower Right Lat, Lon = 35.4°, -115.2°
 Aircraft Heading = 174.9°
 Solar Zenith = 45.8°
 GPS Altitude = 20114. m (MSL)

Citation

Hook, S.J., J.S. Myers, K.J. Thome, M. Fitzgerald, A.B. Kahle, Airborne Sensor Facility NASA Ames Research Center, and R.O. Green. 2022. MASTER: Western Diversity Time Series Campaign, WDTS, California, USA, Spring 2021. ORNL DAAC, Oak Ridge, Tennessee, USA. <https://doi.org/10.3334/ORNLDAAC/1953>

Table of Contents

1. [Dataset Overview](#)
2. [Data Characteristics](#)
3. [Application and Derivation](#)
4. [Quality Assessment](#)
5. [Data Acquisition, Materials, and Methods](#)
6. [Data Access](#)
7. [References](#)
8. [Dataset Revisions](#)

1. Dataset Overview

This dataset includes Level 1B (L1B) and Level 2 (L2) data products from the MODIS/ASTER Airborne Simulator (MASTER) instrument. The spectral data were collected as part of the Western Diversity Time Series (WDTS, formerly HypSPIRI) airborne campaign during nine flights aboard a NASA ER-2 aircraft over selected areas of California and Nevada, U.S., from 2021-02-09 to 2021-04-02. The WDTS campaign will observe California's ecosystems and provide critical information on natural disasters such as volcanoes, wildfires, and drought. MASTER products can identify vegetation type and health and provide a benchmark for the state of the ecosystems against which future changes can be assessed. Data products include L1B georeferenced multispectral imagery of calibrated radiance in 50 bands covering wavelengths of 0.460 to 12.879 micrometers at approximately 50-meter spatial resolution. Derived L2 data products are emissivity in 5 bands in thermal infrared range (8.58 to 12.13 micrometers) and land surface temperature. The L1B file format is HDF-4, and L2 products are provided in ENVI and KMZ formats. In addition, the dataset includes the flight path, spectral band information, instrument configuration, ancillary notes, and summary information for each flight, and browse images derived from each L1B data file.

The MASTER instrument is a modified Daedalus Wildfire scanning spectrometer that flies on a variety of multi-altitude research aircraft and provides spectral information similar to that provided by the Moderate Resolution Imaging Spectroradiometer (MODIS) and the Advanced Spaceborne Thermal Emission and Reflection Radiometer (ASTER), which are aboard two NASA Earth Observing System satellites: Terra and Aqua. The primary goal of this airborne campaign was to demonstrate important science and applications research that is uniquely enabled by the full suite of MASTER thermal infrared bands as well as the contiguous spectroscopic measurements of the AVIRIS (also flown in similar campaigns), or combinations of measurements from both instruments.

Project: [MODIS/ASTER Airborne Simulator](#)

The MODIS/ASTER Airborne Simulator (MASTER) is a scanning spectrometer which flies on a variety of multi-altitude research aircraft and provides data similar to the Moderate Resolution Imaging Spectroradiometer (MODIS) and the Advanced Spaceborne Thermal Emission and Reflection Radiometer (ASTER). MASTER first flew in 1998 and has ongoing deployments as a Facility Instrument in the NASA Airborne Science Program (ASP). MASTER is a joint project involving the Airborne Sensor Facility (ASF) at the Ames Research Center, the Jet Propulsion Laboratory (JPL), and the Earth Resources Observation and Science Center (EROS).

Related Publication

Hook, S.J. Myers, J.J., Thome, K.J., Fitzgerald, M. and A.B. Kahle. 2001. The MODIS/ASTER airborne simulator (MASTER) - a new instrument for earth science studies. *Remote Sensing of Environment* 76:93–102. [https://doi.org/10.1016/S0034-4257\(00\)00195-4](https://doi.org/10.1016/S0034-4257(00)00195-4)

Related Datasets

Additional MASTER datasets are available on the ORNL DAAC [MASTER](#) project page.

Acknowledgments

The MASTER instrument is maintained and operated by the Airborne Sensor Facility (ASF) at NASA Ames Research Center in Mountain View, California, under the oversight of the EOS Project Science Office at NASA Goddard. Data processing was conducted at NASA Ames Research Center and the Jet Propulsion Laboratory at the California Institute of Technology in Pasadena, California.

2. Data Characteristics

Spatial Coverage: Portions of southern California and western Nevada, U.S.

Spatial Resolution: 45 to 50 m

Temporal Coverage: 2020-09-17 to 2020-10-15

Temporal Resolution: One-time estimate

Study Area: Latitude and longitude are given in decimal degrees.

Site	Northernmost Latitude	Southernmost Latitude	Easternmost Longitude	Westernmost Longitude
southern California; western Nevada	40.980	31.951	-112.498	-123.817

Data File Information

This dataset includes a total of 680 data files: 82 files in Hierarchical Data Format (HDF-4; *.hdf) format, 308 ENVI raster files (*.dat and *.hdr) that are compressed (*.zip), 77 files in Keyhole Markup Language Zipped (KMZ; *.kmz) format, 77 Portable Network Graphics (PNG; *.png) files that are compressed (*.zip), 36 text (*.txt) files, 9 archives of text files that are zipped (*.zip), 9 flight maps as GIF (*.gif) images, and 82 browse images in JPEG (*.jpg) format (Table 1).

There are different numbers of each type of file, which corresponds to the number of "flights" and "flight tracks". A "flight" is flown on a single day, and a "flight track" typically refers to a segment of a given flight. The number of flight tracks varies among flights (Table 2).

- There are 9 flights with 82 flight tracks (Table 2).
- For each of 82 flight tracks, there is one L1B data file in HDF format and an auxiliary browse image (*.jpg).
- L2 data are included for 77 of the 82 flight tracks. For each track, there is one L1B data file in KMZ format and five L2 data files in ZIP format.
 - Four of the ZIP files contain L2 ENVI data for emissivity, land surface temperature, geographic coordinates, and quality assurance status. Each ZIP includes a binary data file (*.dat) and its header file (*.hdr).
 - One ZIP file contains three L2 PNG files: RGB composite, single-band emissivity, and land surface temperature.
- For each flight, there is a collection of auxiliary files providing information about the flight and instrument configuration.

The primary data files are named MASTERL1B_#####_CC_YYYYMMDD_EEFF_GGHH_V0J-X.ext (e.g., MASTERL1B_2192800_01_20210209_1937_1945_V01.hdf).

The flight track-level browse images are named MASTERL1B_#####_CC_YYYYMMDD_EEFF_GGHH_V0J.jpg (e.g., MASTERL1B_2192800_01_20210209_1937_1945_V01.jpg).

The deployment-level auxiliary files are named MASTER_#####_YYYYMMDD_X.ext (e.g., MASTER_2192800_20210209_config.txt).

Elements of file names are described as:

= the flight number (see Table 2),
CC = flight track,
YYYYMMDD = date of sampling,
EEFF = starting time at EE hour and FF minute,
GGHH = ending time at GG hour and HH minute,
J = version number for file,
X = the file content (see Table 1), and
ext = "hdf", "gif", "kmz", "jpg", "txt", or "zip", indicating the file extension.

Table 1. File names and descriptions.

File Name	Level	File Type	Total Files	Description
Primary Data Files				
MASTERL1B_#####_CC_YYYYmmDD_EEFF_GGHH_V0J.hdf	L1B	HDF-4	82	Multispectral radiance in 50 bands, pixel coordinates, sensor configuration, aircraft platform data, analysis parameters. The "CalibratedData" variable provides estimates of radiance in units of $W\ m^{-2}\ sr^{-1}$ per micron.
MASTERL1B_#####_CC_YYYYmmDD_EEFF_GGHHV0J-RGB.kmz	L1B	KMZ	77	RGB composite browse image (in KMZ format) derived from corresponding bands of RGB wavelengths of L1B data.
MASTERL2_#####_CC_YYYYmmDD_EEFF_GGHH_V0J-emissivity_tes.zip	L2	ENVI	77	Map of atmospheric corrected emissivity; Temperature and Emissivity Separation (TES) corrected data in 5 bands (wavelengths: 8.58, 9.02, 10.62, 11.32, and 12.13 μm).
MASTERL2_#####_CC_YYYYmmDD_EEFF_GGHH_V0J-images.zip	L2	PNG	77	Three non-georeferenced images depicting (a) RGB composite using selected bands, (b) emissivity from a single band, and (c) land surface temperature.
MASTERL2_#####_CC_YYYYmmDD_EEFF_GGHH_V0J-location.zip	L2	ENVI	77	Latitude and longitude coordinates for pixels in ENVI files.
MASTERL2_#####_CC_YYYYmmDD_EEFF_GGHH_V0J-QAmap.zip	L2	ENVI	77	QA status for each pixel from TES algorithm, where 1 = divergence and 0 = convergence.
MASTERL2_#####_CC_YYYYmmDD_EEFF_GGHH_V0J-surface_temp.zip	L2	ENVI	77	Map of land surface temperature (TES LST) in degrees Kelvin.
Auxiliary Files				
MASTERLAA_#####_CC_YYYYMMDD_EEFF_GGHH_V0J.jpg	L1B	JPEG	82	Browse figures; one image per flight track; multiple tracks per flight.
MASTER_#####_YYYYMMDD_ancillary.txt	-	Text	9	Ancillary information about flight including notes on aircraft platform, mission objective, and data evaluation.
MASTER_#####_YYYYMMDD_config.txt	-	Text	9	Instrument configuration information for flight.
MASTER_#####_YYYYMMDD_flightpath.gif	-	GIF	9	Map showing flight paths.
MASTER_#####_YYYYMMDD_spectral_band_info.txt	-	Text	9	Spectral band information for flight.
MASTER_#####_YYYYMMDD_spectral_response_table.zip	-	Text	9	Spectral response tables by band (ZIP archive of 50 text files).

File Name	Level	File Type	Total Files	Description
-----------	-------	-----------	-------------	-------------

MASTER_BBBBBBBB_YYYYMMDD_summary.txt	-	Text	9	Time and coordinates for start and end of flight tracks along with the number of scan lines, solar and instrument angles, and aircraft altitude. FTLT = flight track number.
--------------------------------------	---	------	---	--

Data File Details

The HDF-4 files contain swath trajectory data using longitude and latitude coordinates. The spatial resolution ranges from 45 m to 50 m and is a function of aircraft altitude.

Table 2. Number of flight tracks for each MASTER flight during this 2021 deployment over California (CA) and Nevada (NV).

Date	Flight Number	Locations (USA)	Flight Tracks	
			Data Level	L2
2021-02-09	2192800	Sierra Nevada Mountains	4	2
2021-02-10	2192900	Santa Barbara Box (Line 1)	11	11
2021-02-24	2193000	Ivanpah Playa (CA/NV)	11	10
2021-03-16	2193200	Lake Tahoe Box (Line 11)	4	4
2021-03-24	2193500	Southern California Box (Line 1)	7	7
2021-03-26	2193600	S.F. Bay Area Box (Line 12)	12	12
2021-03-29	2193700	Yosemite NEON Box (Line 11)	12	12
2021-03-30	2193800	Lake Tahoe Box (Line 11)	12	10
2021-04-02	2193900	Southern California Box (Line 3)	9	9
Total			82	77

3. Application and Derivation

The primary objective of MASTER is to: (a) collect ASTER-like and MODIS-like land datasets to support the validation of the ASTER and MODIS geophysical retrieval algorithms; (b) collect these datasets at a higher resolution than the spaceborne datasets to permit scaling studies and comparisons with in-situ measurements; and (c) under fly the EOS-AM1 ASTER and MODIS sensors to provide an additional radiometric calibration to assist with in-flight instrument performance characterization. Calibration is particularly important for ASTER where on-board calibration is dependent on a single black body in the TIR and only partial aperture illumination in the VNIR.

A secondary objective of MASTER is to: (a) provide both a backup instrument and backup modules for the current MODIS Airborne simulator, which is committed to a program of atmospheric and oceanic measurements; and (b) provide a wider spectral and dynamic range alternative to the use of the Thematic Mapper (TM) airborne simulator and Thermal Infrared Multispectral Scanner (TIMS) airborne scanners (JPL, 2021b).

MASTER imagery has been used for mapping wildfires and their impacts (Veraverbeke et al., 2011), land cover (Li and Moon, 2004), coral reefs (Capolsini et al., 2003), and urban heat islands (Zhao and Wentz, 2016).

4. Quality Assessment

The MASTER instrument channels are calibrated spectrally and radiometrically in the laboratory preflight and postflight. The mid-infrared and thermal infrared channels (26–50) are also radiometrically calibrated in-flight by viewing an internal hot and cold blackbody with each scanline (Hook et al., 2001). Three calibration and validation experiments were conducted in 1998–2001 (Hook et al., 2001; JPL, 2021a). Spectral response information for this deployment is included in the files named MASTER_BBBBBBBB_YYYYMMDD_spectral_response_table.zip.

5. Data Acquisition, Materials, and Methods

The MASTER instrument was developed by the NASA Ames Research Center in conjunction with the Jet Propulsion Laboratory. The instrument consists of three key components: the scanning spectrometer, the digitizer, and the storage system. The scanning unit was built by Sensys Technology (formerly Daedalus Enterprises) and the digitizer was a collaborative effort between Berkeley Camera Engineering and the Ames Airborne Sensor Facility (ASF, 2021). The data storage system and overall system integration were also provided by the ASF.

The MASTER instrument is similar to the MODIS Airborne Simulator (MAS) developed by the MODIS project (King et al., 1996). However, it has two key differences. First, MASTER supports a variety of scan speeds allowing it to acquire contiguous imagery from a variety of altitudes with different pixel sizes. Second, the channel positions are configured to closely match those of ASTER and MODIS. A detailed description of the instrument and optical system are provided by Hook et al. (2001) and King et al. (1996), respectively.

The Western Diversity Time Series (WDTS, formerly HyspIRI) program will observe California's ecosystems and provide critical information on natural disasters such as volcanoes, wildfires, and drought. It will provide a benchmark on the state of the ecosystems against which future changes can be assessed, as the instruments will be capable of identifying vegetation type and health. The WDTS Airborne Campaign is a multi-year effort to collect seasonal VIS-SWIR and TIR airborne scanner data using both AVIRIS and MASTER remote sensing instruments aboard the ER-2 high-altitude platform (ASF, 2021).

For this campaign, the MASTER instrument was flown on NASA's ER-2 aircraft at altitudes of 17,697–20,124 m above sea level.




Figure 2. Flight tracks over California and Nevada represented as rectangular polygons. Basemap: © OpenStreetMap contributors.

The L2 data are derived from the L1B files, and the primary L2 products are emissivity in five bands (wavelengths: 8.58, 9.02, 10.62, 11.32, and 12.13 μm) and land surface temperature (LST). Emissivity and LST were corrected using a Temperature and Emissivity Separation (TES) algorithm (Coll et al., 2001).




Figure 3. Flight path for Flight 2192900, flown on 10 February 2021. Flight 2192900 and 11 flight tracks occurred over southern California, U.S., in the vicinity of Santa Barbara and the Channel Islands. Source: MASTER_2192900_20210210_flightpath.gif

6. Data Access

These data are available through the Oak Ridge National Laboratory (ORNL) Distributed Active Archive Center (DAAC).

[MASTER: Western Diversity Time Series Campaign, WDTS, California, USA, Spring 2021](#)

Contact for Data Center Access Information:

- E-mail: uso@daac.ornl.gov
- Telephone: +1 (865) 241-3952

7. References

- ASF. 2021. Campaign summary information: HypSPIRI / WDTS Airborne Campaign. Airborne Sensor Facility, Airborne Science Program, NASA Ames Research Center, Moffett Field, California. https://asapdata.arc.nasa.gov/sensors/master/data/deploy_html/hyspiri_home.html
- Capolsini, P., S. Andréfouët, C. Rion, and C. Payri. 2003. A comparison of Landsat ETM+, SPOT HRV, Ikonos, ASTER, and airborne MASTER data for coral reef habitat mapping in South Pacific islands. *Canadian J. Remote Sensing* 29:187-200. <https://doi.org/10.5589/m02-088>
- Coll, C., V. Caselles, E. Rubio, F. Sospedra, and E. Valor. 2001. Temperature and emissivity separation from calibrated data of the Digital Airborne Imaging Spectrometer. *Remote Sensing of Environment* 76:250-259. [https://doi.org/10.1016/S0034-4257\(00\)00207-8](https://doi.org/10.1016/S0034-4257(00)00207-8)
- Hook, S.J. Myers, J.J., Thome, K.J., Fitzgerald, M., and A.B. Kahle. 2001. The MODIS/ASTER airborne simulator (MASTER) - a new instrument for earth science studies. *Remote Sensing of Environment* 76:93-102. [https://doi.org/10.1016/S0034-4257\(00\)00195-4](https://doi.org/10.1016/S0034-4257(00)00195-4)
- JPL. 2021a. Calibration and Validation, MASTER: MODIS/ASTER Airborne Simulator. Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, USA. <https://masterprojects.jpl.nasa.gov/cal-val>
- JPL. 2021b. Science objectives, MASTER: MODIS/ASTER Airborne Simulator. Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, USA. <https://masterprojects.jpl.nasa.gov/objectives>
- King, M.D., W.P. Menzel, P.S. Grant, J.S. Myers, G.T. Arnold, S.E. Platnick, L.E. Gumley, S.C. Tsay, C.C. Moeller, M. Fitzgerald, K.S. Brown, and F.G. Osterwisch. 1996. Airborne scanning spectrometer for remote sensing of cloud, aerosol, water vapor and surface properties. *J. Atmospheric and Oceanic Technology* 13:777-794. [https://doi.org/10.1175/1520-0426\(1996\)013<0777:ASSFRS>2.0.CO;2](https://doi.org/10.1175/1520-0426(1996)013<0777:ASSFRS>2.0.CO;2)
- Li, P., and W.M. Moon. 2004. Land cover classification using MODIS-ASTER airborne simulator (MASTER) data and NDVI: A case study of the Kochang area, Korea. *Canadian J. Remote Sensing* 30:123-126. <https://doi.org/10.5589/m03-061>
- Veraverbeke, S., S. Harris, and S. Hook. 2011. Evaluating spectral indices for burned area discrimination using MODIS/ASTER (MASTER) airborne simulator data. *Remote Sensing of Environment* 115:2702-2709. <https://doi.org/10.1016/j.rse.2011.06.010>
- Zhao, Q., and E.A. Wentz. 2016. A MODIS/ASTER Airborne Simulator (MASTER) imagery for urban heat island research. *Data* 1:7. <https://doi.org/10.3390/data1010007>

8. Dataset Revisions

Version	Release Date	Revision Notes
1.1	2023-01-27	Level 2 files added to dataset. User Guide updated.
1.0	2022-05-12	Original publication



[Privacy Policy](#) | [Feedback](#) | [Help](#)



Home

About Us

Mission
Data Use and Citation
Policy
User Working Group
Partners

Get Data

Science Themes
NASA Projects
All Datasets

Submit Data

Submit Data Form
Data Scope and
Acceptance
Data Authorship Policy
Data Publication Timeline
Detailed Submission
Guidelines

Tools

MODIS
THREDDS
SDAT
Daymet
Airborne Data Visualizer
Soil Moisture Visualizer
Land - Water Checker

Resources

Learning
Data Management
News
Earthdata Forum [↗](#)

Contact Us

