

NASA Technical Memorandum 4153

**Global Data Bases on Distribution,
Characteristics and Methane
Emission of Natural Wetlands:
Documentation of Archived Data Tape**

Elaine Matthews
Goddard Institute for Space Studies
New York, New York



National Aeronautics and
Space Administration

Office of Management

Scientific and Technical
Information Division

1989

INTRODUCTION

Global digital data bases on the distribution and environmental characteristics of natural wetlands, compiled by Matthews and Fung (1987), have been archived on tape. These data bases were developed to evaluate the role of wetlands in the annual emission of methane from terrestrial sources. Five global 1° latitude by 1° longitude arrays are included on the tape. They are: (1) wetland data source, (2) wetland type, (3) fractional inundation, (4) vegetation type and (5) soil type. The first three data bases on wetland locations were published by Matthews and Fung (1987). The last two arrays contain ancillary information about these wetland locations: vegetation type from the data of Matthews (1983) and soil type from the data of Zobler (1986). This short paper is designed only to document the tape, and briefly explain the data sets and their initial application to estimating the annual emission of methane from natural wetlands. For complete discussions of the data, including sources and uncertainties, consult original publications listed in the references.

Globally, wetlands occur in a total of 3233 1° cells and occupy about $5.3 \times 10^{12} \text{m}^2$ (Matthews and Fung 1987). Each of the five data bases here includes information for the 3233 wetland locations; cells of other (non-wet) land and water are masked with unique values.

The outline below gives information about array characteristics such as dimensions, read formats, record lengths, block sizes and value ranges. Sections 1–5 following the outline contain descriptions and translation tables of the individual data bases. The final section includes information on the first calculation of global methane emission from natural wetlands done by Matthews and Fung (1987) using these data sets in conjunction with field measurements of methane fluxes from various ecosystems and simple latitudinal estimates about the seasonality of methane production.

References

FAO (1971–1981): Soil Map of the World, Vols. 1–10 (1:5M scale maps and accompanying texts), UNESCO, Paris.

Matthews, E. (1983): Global vegetation and land use: new high-resolution data bases for climate studies. J. Clim. Appl. Meteorol., 22, 474–487.

Matthews, E. and I. Fung (1987): Methane emission from natural wetlands: global distribution, area and environmental characteristics of sources. Global Biogeochem. Cycles, 1, 61-86.

UNESCO (1973): International classification and mapping of vegetation. UNESCO, Paris.

Zobler, L. (1986): A world soil file for global climate modeling. NASA Technical Memorandum 87802.

OUTLINE

Name	Size	Type	Format	Description
<u>1. DATA SOURCES</u>				
ISRC	(360,180)	Integer*4	do 10 j=1,180 read(20,'(250I4,110I4)') *(ISRC(i,j),i=1,360) 10 continue	wetland: 1-7 other land: 0 water: -1 see Section 1, Table 1
<u>2. WETLAND TYPES</u>				
IWET	(360,180)	Integer*4	do 10 j=1,180 read(20,'(250I4,110I4)') *(IWET(i,j), I=1,360) 10 continue	wetland: 1-12 other land: 0 water: -1 see Section 2, Table 2
<u>3. FRACTIONAL INUNDATION</u>				
FRIN	(360,180)	Real*4	do 10 j=1,180 read(20,'(250F4.0,110F4.0)') *(FRIN(i,j), I=1,360) 10 continue	wetland: 1.-100. other land: 0. water: -1. see Section 3, Table 3
<u>4. VEGETATION TYPES</u>				
IVEG	(360,180)	Integer*4	do 10 j=1,180 read(20,'(250I4,110I4)') *(IVEG(i,j), I=1,360) 10 continue	wetland: 1-178 other land: 0 water: -1 see Section 4, Table 4
<u>5. SOIL TYPES</u>				
ISOL	(360,180)	Integer*4	do 10 j=1,180 read(20,'(250I4,110I4)') *(ISOL(i,j), i=1,360) 10 continue	wetland: 1-107 other land: 0 water: -1 see Section 5, Table 5

The tape: 9 track, 1600 bpi, ASCII, BLP, RECFM=F

The arrays: record length = 1440, blocksize = 1440.
 (i,j) arrays are (360,180), 1° (lon,lat) resolution
 j = 1,180: j = 1: 1° band from 90°S to 89°S ...
 j = 180: 1° band from 89°N to 90°N
 i = 1,360: i = 1: 1° band from 180° (dateline) to 179°W ...
 i = 360: 1° band from 179°E to 180° (dateline)

Section 1. DATA SOURCES

Description: ISRC

The array of data sources (ISRC) gives information on the source or combination of sources for wetland designations. The three independent data sources integrated to produce the final data base are: (1) vegetation (Matthews 1983) classified with the UNESCO (1973) system (reference UNESCO in Table 1); (2) ponded soils from the data base of Zobler (1986) based on FAO soil maps (reference FAO); and (3) fractional inundation compiled specifically for the Matthews and Fung (1987) study of methane emission from wetlands based on a global series of 1:1M scale Operation Navigation Charts (reference ONC). All cells identified as wetland by any single source or combination of sources are designated as wetlands in the integrated data set given here. The three data sources provide a total of seven source combinations as shown in Table 1. For example, type 1 locations are targeted as wetlands by all three source data bases while type 6 locations are identified as wetlands only by the ONC inundation data base.

Table 1. ISRC: Data Sources For Wetland Locations

Tape	M&F ¹	Data Source Combinations
1	A	UNESCO + FAO + ONC
2	B	UNESCO + ONC
3	C	UNESCO + FAO
4	D	UNESCO
5	E	ONC + FAO
6	F	ONC
7	G	FAO
0		other (non-wet) land
-1		water

¹ letter reference from Figure 5 of Matthews and Fung (1987)

Section 2. WETLAND TYPE

Description: IWET

A total of twelve wetland types are identified in the array IWET. Twenty-eight wetland vegetation types were originally chosen from the UNESCO system and classified into five major wetland groups on the basis of vegetational features and environmental characteristics that affect methane emission (as discussed in Matthews and Fung 1987); these are: (1) forested bog, (2) nonforested bog, (3) forested swamp, (4) nonforested swamp and (5) alluvial formations (see Table 2). Many additional locations identified as wetlands from the FAO and ONC data bases were occupied by about 100 vegetation types other than these 28 explicit wetland ecosystems. For the calculation of methane emission, vegetation types in these additional cells were initially grouped into seven categories on the basis of climate and vegetational structure (6A through 6G under G1, Table 2); each category was then associated with one of the five major wetland groups. Column G2 (Table 2) shows the secondary groupings of the additional seven vegetation categories with five major wetland groups. To allow for alternative combinations, all twelve wetland types are identified separately in the archived data base, as indicated in the Tape column.

Table 2. IWET: Wetland Types and Wetland Groupings

Tape	G1 ¹	G2 ²	Description
1	1	1	forested bog
2	2	2	nonforested bog
3	3	3	forested swamp
4	4	4	nonforested swamp
5	5	5	alluvial formations
6	6A	3	tropical/subtropical forest/woodland
7	6B	3	temperate forest/woodland
8	6C	1	high-latitude temperate/boreal forest/woodland/shrub
9	6D	3	shrubland; xeromorphic formations; desert
10	6E	4	wooded grassland
11	6F	4	nonwooded grassland
12	6G	2	tundra
0			other (non-wet) land
-1			water

¹ type number from Table 2b and Table 5 in Matthews and Fung (1987)

² wetland group with which type was associated for methane emission calculation (Matthews and Fung, 1987)

Section 3. FRACTIONAL INUNDATION

Description: FRIN

The fractional inundation data base (FRIN) was compiled from a series of 1:1M scale Operational Navigation Charts (ONC). The values, which give the inundated proportion of 10^6 cells, were used to calculate the global wetland area of $5263 \times 10^9 \text{m}^2$ (Matthews and Fung, 1987). For cells targeted as wetlands by UNESCO vegetation or FAO ponding but not by the ONC inundation data base (i.e., cases 3, 4 and 7 in Table 1), the fractional wetland coverage was prescribed as the mean inundation for the vegetation type occupying the cell; these locations were incorporated into the fractional inundation data base.

Table 3. FRIN: Fractional Inundation

Tape	Description
1. - 100. ¹	wetland
0.	other (non-wet) land
-1.	water

¹ unit is percent

Section 4. VEGETATION TYPE

Description: IVEG

The vegetation type array identifies UNESCO (1973) vegetation codes of wetland sites from the data set of Matthews (1983). Table 4 is a list of the integers in IVEG (Tape column), their associated vegetation codes and a brief description of the vegetation types. The UNESCO classification system is hierarchical, with a maximum of five alpha-numeric components in each entry. Table 4 gives the complete series of vegetation codes that occur in the full vegetation data base compiled by Matthews (1983) (i.e., codes accompanied by integers in the Tape column of the table) although not all occur in the wetlands data base of Matthews and Fung (1987) given here. Codes unaccompanied by integers occur in neither the vegetation nor wetlands data bases but are included for clarity in following the classification structure.

Table 4. IVEG: UNESCO Vegetation Types of Wetland Locations

Tape	Code	Description
	1.A	EVERGREEN FOREST
1	1.A.1	Tropical evergreen rainforest
2	1.A.1a	lowland
3	1.A.1b	submontane
4	1.A.1c	montane
5	1.A.1c2	needleleaved
6	1.A.1e	cloud
7	1.A.1f	alluvial
8	1.A.1f1	frequently flooded.
9	1.A.1f3	seasonally water-logged
10	1.A.1g	swamp
11	1.A.1g2	dominated by palms
12	1.A.1h	bog
13	1.A.2	Tropical/subtropical evergreen seasonal forest
14	1.A.2a	lowland
15	1.A.2b	submontane
16	1.A.2b2	needleleaved
17	1.A.2c	montane
18	1.A.3	Tropical/subtropical semi-deciduous forest
19	1.A.4	Subtropical evergreen rainforest
20	1.A.4b	submontane
21	1.A.4c	montane
22	1.A.4c2	needleleaved
23	1.A.4f	alluvial
24	1.A.5	Mangrove forest
25	1.A.6	Temperate/subpolar evergreen rainforest
26	1.A.6a	temperate
27	1.A.6a1	broadleaved
28	1.A.6a2	broadleaved with needleleaved trees
29	1.A.7	Temperate evergreen broadleaved seasonal forest
30	1.A.8	Evergreen broadleaved sclerophyllous forest (winter rain)
31	1.A.8a	lowland/submontane
32	1.A.8b	lowland/submontane, generally less than 50 m tall
33	1.A.9	Tropical/subtropical evergreen needleleaved forest
34	1.A.9a	lowland/submontane
35	1.A.9b	montane/subalpine
36	1.A.10	Temperate/subpolar evergreen needleleaved forest
37	1.A.10c	with conical crowns
38	1.A.10d	with cylindro-conical crowns (boreal)
39	1.A.10e	with cylindro-conical crowns (boreal): water-logged

- 1.B DECIDUOUS FOREST**
- 40 1.B.1 Tropical/subtropical drought–deciduous forest
- 41 1.B.1a broadleaved lowland/submontane
- 42 1.B.1b montane (and cloud)
- 43 1.B.2 Cold–deciduous broadleaved forest with evergreen trees or shrubs
- 44 1.B.2a with evergreen broadleaved trees and climbers
- 45 1.B.2b with evergreen needleleaved trees
- 46 1.B.2c subalpine and subpolar
- 47 1.B.2d subalpine/subpolar alluvial
- 48 1.B.2e waterlogged
- 49 1.B.3 Cold–deciduous forest without evergreen trees
- 50 1.B.3a temperate lowland/submontane
- 51 1.B.3b montane/boreal
- 52 1.B.3b1 broadleaved
- 53 1.B.3b2 needleleaved (e.g. Larix)
- 54 1.B.3c subalpine/subpolar
- 55 1.B.3d alluvial
- 56 1.B.3d2 regularly flooded with abundant herbaceous undergrowth
- 57 1.B.3e swamp or peat
- 1.C EXTREMELY XEROMORPHIC FOREST**
- 58 1.C.1 Extremely xeromorphic sclerophyllous–dominated forest
- 59 1.C.2 Extremely xeromorphic thorn forest
- 60 1.C.2a mixed deciduous–evergreen
- 61 1.C.2b deciduous
- 62 1.C.2c evergreen
- 2.A EVERGREEN WOODLAND**
- 63 2.A.1 Evergreen broadleaved woodland
- 64 2.A.2 Evergreen needleleaved woodland
- 65 2.A.2a with rounded crowns
- 66 2.A.2a1 with evergreen sclerophyllous understory (Mediterranean)
- 67 2.A.2b with conical crowns (subalpine)
- 68 2.A.2c with cylindro–conical crowns (boreal)
- 69 2.A.2d waterlogged
- 2.B DECIDUOUS WOODLAND**
- 70 2.B.1 Tropical/subtropical drought–deciduous woodland
- 71 2.B.1a lowland/submontane, broadleaved
- 72 2.B.1b montane (and cloud)
- 73 2.B.2 Cold–deciduous woodland with evergreen trees
- 2.B.3 Cold–deciduous woodland without evergreen trees
- 74 2.B.3a broadleaved
- 75 2.B.3b needleleaved
- 76 2.B.3b2 mixed broadleaved–needleleaved
- 2.C EXTREMELY XEROMORPHIC WOODLAND**
- 77 2.C Extremely xeromorphic woodland
- 78 2.C.1 Extremely xeromorphic sclerophyllous–dominated woodland
- 79 2.C.2 Extremely xeromorphic thorn woodland
- 80 2.C.2a mixed deciduous–evergreen
- 81 2.C.2c deciduous
- 82 2.C.3 Extremely xeromorphic succulent woodland

- 3.A EVERGREEN SHRUBLAND**
- 83 3.A.1 Evergreen broadleaved shrubland or thicket
84 3.A.1a low bamboo thicket
85 3.A.1d sclerophyllous shrubland or thicket
86 3.A.2 Evergreen needleleaved or microphyllous shrubland or thicket
87 3.A.2a needleleaved
88 3.A.3b microphyllous
- 3.B DECIDUOUS SHRUBLAND**
- 89 3.B.1 Drought-deciduous shrubland with evergreens
90 3.B.2 Drought-deciduous shrubland without evergreens
91 3.B.2b subalpine/subpolar
92 3.B.3 Cold-deciduous shrubland
93 3.B.3b subalpine/subpolar
93 3.B.3b1 dwarf shrubland, with forbs
94 3.B.3b2 dwarf shrubland, with lichens
95 3.B.3c alluvial
- 3.C EXTREMELY XEROMORPHIC SUBDESERT SHRUBLAND**
- 96 3.C Extremely xeromorphic subdesert shrubland
97 3.C.1 Extremely xeromorphic evergreen subdesert shrubland
98 3.C.1a evergreen
99 3.C.1a1 broadleaved
100 3.C.1a2 microphyllous, or leafless with green stems
101 3.C.1a3 succulent
102 3.C.1b semi-deciduous
103 3.C.1b1 facultatively deciduous
104 3.C.2 Extremely xeromorphic deciduous subdesert shrubland
105 3.C.2b with succulents
- 4.A DWARF SHRUBLAND**
- 106 4.A.1 Evergreen dwarf-shrub thicket
107 4.A.2 Evergreen dwarf shrubland
108 4.A.2a dense cushion
109 4.A.3 Mixed evergreen dwarf shrub/herbaceous formation
- 4.C EXTREMELY XEROMORPHIC DWARF SHRUBLAND**
- 110 4.C Extremely xeromorphic subdesert dwarf shrubland
110 4.C.1 Extremely xeromorphic subdesert dwarf shrubland
111 4.C.1a evergreen
112 4.C.2 Extremely xeromorphic deciduous subdesert dwarf shrubland
- 4.D TUNDRA**
- 113 4.D Tundra
114 4.D.1 Mainly bryophyte tundra
115 4.D.2 Mainly lichen tundra
116 4.D.2a with caespitose dwarf shrubs and moss
117 4.D.2b with creeping or matted dwarf shrubs and moss
- 4.E MOSSY BOGS**
- 118 4.E Mossy bog formations with dwarf shrubs
118 4.E.2 Non-raised mossy bog
119 4.E.2b string bog

- 5.A TALL GRASSLAND**
- 120 **5.A.1 Tall grassland with 10–40% tree cover**
 121 5.A.1a with evergreen broadleaved tree cover
 122 5.A.1a1 wet or flooded most of year
 123 5.A.1c with deciduous broadleaved tree cover
 124 5.A.1c1 seasonally flooded
 125 5.A.1c2 with deciduous broadleaved tree cover
 126 **5.A.2 Tall grassland with <10% tree cover**
 127 5.A.2c with deciduous broadleaved tree cover
 128 5.A.3c with deciduous broadleaved shrub cover
 129 **5.A.4 Tall grassland with tuft plant cover (usually palms)**
 130 **5.A.5 Tall grassland without woody cover**
 131 5.A.5a tropical grassland
 132 5.A.5a1 seasonally flooded
 133 5.A.5a2 wet or flooded most of year
- 5.B MEDIUM GRASSLAND**
- 134 **5.B.1 Medium grassland with 10–40% tree cover**
 135 5.B.1a with evergreen broadleaved tree cover
 136 5.B.1a1 wet or flooded most of year
 137 5.B.1b with semi-evergreen broadleaved tree cover
 138 5.B.1c with deciduous broadleaved tree cover
 139 **5.B.2 Medium grassland with <10% tree cover**
 140 **5.B.3 Medium grassland with shrub cover**
 141 5.B.3c with deciduous broadleaved shrub cover
 142 5.B.3e with deciduous thorny shrub cover
 143 **5.B.4 Medium grassland with open cover of tuft plants (usually palms)**
 144 5.B.4a subtropical, with open groves of palms
 145 **5.B.5 Medium grassland without woody cover**
 5.B.5a mainly sod grasses
 146 5.B.5a1 wet or flooded most of year
 147 5.B.5a2 on sandy soil or dunes
 148 5.B.5b mainly bunch grasses
 149 5.B.5b2 wet or flooded most of year
- 5.C SHORT GRASSLAND**
- 150 **5.C.1 Short grassland with 10–40% tree cover**
 151 5.C.1a with evergreen broadleaved tree cover
 152 5.C.1c with deciduous broadleaved tree cover
 153 5.C.1d with evergreen needleleaved tree cover
 154 **5.C.2 Short grassland with <10% tree cover**
 5.C.2a with evergreen broadleaved tree cover
 155 5.C.2a1 seasonally flooded
 156 5.C.2c with deciduous broadleaved tree cover
 157 **5.C.3 Short grassland with shrub cover**
 158 5.C.3b with semi-evergreen broadleaved shrub cover
 159 5.C.3c with deciduous broadleaved shrub cover
 160 5.C.3e with deciduous thorny shrub cover
 161 **5.C.5 Short grassland without woody cover**
 162 5.C.5a tropical alpine, open/closed bunch-grasses with tuft-plant cover
 163 5.C.5b tropical alpine, open bunch grasses
 164 5.C.5d bunch grasses of varying coverage with dwarf shrubs
 165 **5.C.6 Short grassland without woody cover**
 166 5.C.6a short-grass communities in semi-arid climates
 167 5.C.6b bunch-grass communities (tussock)

- 168 5.C.7 **Short to medium tall mesophytic grassland (meadow)**
- 169 5.C.7a sodgrass communities, forbs in low altitude, cool humid climates
- 170 5.C.7b alpine/subalpine meadows, high latitudes
- 171 5.C.7b2 alpine/subalpine meadows, high latitudes, rich in dwarf shrubs
- 172 5.C.7b3 snow-bed communities in high latitude alpine/subalpine meadows
- 173 5.C.8 **Graminoid tundra**
- 174 5.C.8a bunch-form with mosses and lichens (*Eriophorum*)

5.D FORB FORMATIONS

- 175 5.D.2 **Low forb communities (<1m)**
- 176 5.D.2a perennial flowering forbs and ferns

OTHER

- 177 6 desert
 - 178 7 ice
 - 0 other non-wet land
 - 1 water
-

Section 5. SOIL TYPE

Description: ISOL

Soil types for wetland sites (ISOL) are from the data of Zobler (1986), digitized from the global series of soil maps produced by FAO (1971–1981). Locations are identified by integers associated with 106 soil units; ice is identified as 107. Table 5 lists integer values on the tape, along with FAO codes and FAO soil unit names. Note that FAO soil unit data in this array are independent of the distribution of ponded phase soils (from Zobler 1986) selected as one of three sources used to identify wetland locations by Matthews and Fung (1987).

Table 5. ISOL: FAO Soil Types of Wetland Locations

Tape	Code	Name	Tape	Code	Name
	A	<u>ACRISOL</u>		G	<u>GLEYSOL</u>
1	AF	Ferric Acrisol	29	GC	Calcaric Gleysol
2	AG	Gleyic Acrisol	30	GD	Dystric Gleysol
3	AH	Humic Acrisol	31	GE	Eutric Gleysol
4	AO	Orthic Acrisol	32	GH	Humic Gleysol
5	AP	Plinthic Acrisol	33	GM	Mollic Gleysol
			34	GP	Plinthic Gleysol
			35	GX	Gelic Gleysol
	B	<u>CAMBISOL</u>		H	<u>PHAEOZEM</u>
6	BC	Chromic Cambisol			
7	BD	Dystric Cambisol			
8	BE	Eutric Cambisol	36	HC	Calcaric Phaeozem
9	BF	Ferralic Cambisol	37	HG	Gleyic Phaeozem
10	BG	Gleyic Cambisol	38	HH	Haplic Phaeozem
11	BH	Humic Cambisol	39	HL	Luvic Phaeozem
12	BK	Calcic Cambisol			
13	BV	Vertic Cambisol			
14	BX	Gelic Cambisol	40	I	<u>LITHOSOL</u>
	C	<u>CHERNOZEM</u>		J	<u>FLUVISOL</u>
15	CG	Glossic Chernozem	41	JC	Calcaric Fluvisol
16	CH	Haplic Chernozem	42	JD	Dystric Fluvisol
17	CK	Calcic Chernozem	43	JE	Eutric Fluvisol
18	CL	Luvic Chernozem	44	JT	Thionic Fluvisol
	D	<u>PODZOLUVISOL</u>		K	<u>KASTANOZEM</u>
19	DD	Dystric Podzoluvisol	45	KH	Haplic Kastanozem
20	DE	Eutric Podzoluvisol	46	KK	Calcic Kastanozem
21	DG	Gleyic Podzoluvisol	47	KL	Luvic Kastanozem
22	E	<u>RENDZINA</u>		L	<u>LUVISOL</u>
			48	LA	Albic Luvisol
	F	<u>FERRALSOL</u>	49	LC	Chromic Luvisol
23	FA	Acric Ferralsol	50	LF	Ferric Luvisol
24	FH	Humic Ferralsol	51	LG	Gleyic Luvisol
25	FO	Orthic Ferralsol	52	LK	Calcic Luvisol
26	FP	Plinthic Ferralsol	53	LO	Orthic Luvisol
27	FR	Rhodic Ferralsol	54	LP	Plinthic Luvisol
28	FX	Xanthic Ferralsol	55	LV	Vertic Luvisol

	M	<u>GREYZEM</u>		T	<u>ANDOSOL</u>
56	MG	Gleyic Greyzem	81	TH	Humic Andosol
57	MO	Orthic Greyzem	82	TM	Mollic Andosol
			83	TO	Ochric Andosol
			84	TV	Vitric Andosol
	N	<u>NITOSOL</u>			
58	ND	Dystric Nitosol	85	U	<u>RANKER</u>
59	NE	Eutric Nitosol			
59	NH	Humic Nitosol			
60				V	<u>VERTISOL</u>
	O	<u>HISTOSOL</u>	86	VC	Chromic Vertisol
			87	VP	Pellic Vertisol
61	OD	Dystric Histosol			
62	OE	Eutric Histosol			
63	OX	Gelic Histosol		W	<u>PLANOSOL</u>
	P	<u>PODZOL</u>	88	WD	Dystric Planosol
64	PF	Ferric Podzol	89	WE	Eutric Planosol
65	PG	Gleyic Podzol	90	WH	Humic Planosol
66	PH	Humic Podzol	91	WM	Mollic Planosol
67	PL	Leptic Podzol	92	WS	Solodic Planosol
68	PO	Orthic Podzol	93	WX	Gelic Planosol
69	PP	Placic Podzol			
	Q	<u>ARENOSOL</u>		X	<u>XEROSOL</u>
70	QA	Albic Arenosol	94	XH	Haplic Xerosol
71	QC	Cambic Arenosol	95	XK	Calcic Xerosol
72	QF	Ferralic Arenosol	96	XL	Luvic Xerosol
73	QL	Luvic Arenosol	97	XY	Gypsic Xerosol
	R	<u>REGOSOL</u>		Y	<u>YERMOSOL</u>
			98	YH	Haplic Yermosol
			99	YK	Calcic Yermosol
74	RC	Calcaric Regosol	100	YL	Luvic Yermosol
75	RD	Dystric Regosol	101	YT	Takyrlic Yermosol
76	RE	Eutric Regosol	102	YY	Gypsic Yermosol
77	RX	Gelic Regosol			
	S	<u>SOLONETZ</u>		Z	<u>SOLONCHAK</u>
			103	ZG	Gleyic Solonchak
78	SG	Gleyic Solonetz	104	ZM	Mollic Solonchak
79	SM	Mollic Solonetz	105	ZO	Orthic Solonchak
80	SO	Orthic Solonetz	106	ZT	Takyrlic Solonchak
			107		ice
			0		other (non-wet) land
			-1		water

SECTION 6. METHANE EMISSION FROM NATURAL WETLANDS

The data bases presented here were developed to determine the distribution and characteristics of natural wetlands on a global scale and to evaluate the role of wetlands in the annual emission of methane from terrestrial sources. Table 6 (from Matthews and Fung, 1987) shows the latitudinal distribution of wetland areas and the annual methane emission from wetland groups (see Section 2 and Table 2 for explanation). The length of methane emission periods was derived from simple assumptions about the likely duration of thaw conditions in high latitudes and inundation conditions in the subtropics and tropics.

The latitudinal distribution of wetland area derived here shows approximately 50% of the world's wetlands, primarily peat-rich bogs, concentrated in the zone between 50°N and 70°N. Peat-poor swamps, with lower flux rates and longer methane production seasons, are prevalent in low latitudes; about one-third of the wetlands are spread throughout the area from 10°N to 30°S. In this study, the distribution of emission generally parallels that of areas since large daily flux rates for short seasons in the high latitudes balance lower fluxes for longer periods in the tropics and subtropics. The total annual emission of methane from natural wetlands was estimated to be ~110 Tg (10^{12} g), equal to about 20% of the total terrestrial source strength.

Wetlands are generally acknowledged to play a significant role in the global methane budget. The series of data sets presented here, integrated into a comprehensive data base of wetlands and their attributes, provides a framework to analyze and evaluate the role of wetlands in the methane cycle. However, estimates of the magnitude and seasonality of this source may be considered preliminary; further constraints on the wetland source will be provided by measurements in a greater variety of representative wetland ecosystems and throughout methane production seasons and by isotopic measurements.

Table 6. Latitudinal Distribution of Wetland Areas and the Annual Methane Emission From Wetland Groups
(from Matthews and Fung, 1987)

Group	Forested Bog		Nonforested Bog		Forested Swamp			Nonforested Swamp			Alluvial	Total		
	1	6C	2	6G	3	6A	6B	6D	4	6E	6F		5	
Emission Rate, $\text{gCH}_4\text{m}^{-2}\text{d}^{-1}$	0.2		0.2		0.07			0.12			0.03			
<i>Latitude</i>	<i>Period, days</i>	<i>Area x 10⁹m²</i>												
90°N - 80°N	100	-	-	-	-	-	-	-	-	-	-	-	-	-
80°N - 70°N	100	-	5	53	48	-	-	-	7	-	-	-	-	112
70°N - 60°N	100	35	889	334	71	-	-	-	-	-	26	-	-	1355
60°N - 50°N	150	238	556	369	9	22	-	-	1	-	30	10	-	1235
50°N - 40°N	150	95	96	1	0	5	-	0	25	19	64	11	-	319
40°N - 30°N	150	-	53	-	-	-	0	6	25	-	10	25	9	128
30°N - 20°N	180	-	25	-	-	-	11	2	23	16	7	9	1	94
20°N - 10°N	180	-	-	-	-	13	61	-	14	59	129	-	-	276
10°N - 0°	180	16	-	12	-	64	139	0	7	76	79	9	28	431
0° - 10°S	180	64	-	-	-	69	213	6	10	8	49	4	62	484
10°S - 20°S	180	-	-	-	-	3	99	10	17	77	107	25	23	360
20°S - 30°S	180	-	-	-	-	-	21	-	159	40	60	14	39	333
30°S - 40°S	150	5	-	-	-	-	-	18	33	11	10	23	33	132
40°S - 50°S	150	-	-	-	-	-	-	-	3	-	-	-	-	3
50°S - 60°S	150	-	-	-	-	-	-	-	-	-	-	-	-	-
Total Area		453	1624	769	128	177	545	43	323	305	572	130	195	5263
		<i>Emission x 10¹²g</i>												
90°N - 80°N		-	-	-	-	-	-	-	-	-	-	-	-	-
80°N - 70°N		-	0.1	1.1	1.0	-	-	-	0.1	-	-	-	-	2.3
70°N - 60°N		0.7	17.8	6.7	1.4	-	-	-	-	-	0.3	-	-	26.9
60°N - 50°N		7.1	16.7	11.1	0.3	0.2	-	-	0	-	0.6	0.2	-	36.2
50°N - 40°N		2.9	2.9	0	0	0.1	-	0	0.3	0.3	1.2	0.2	-	7.9
40°N - 30°N		-	1.6	-	-	-	0	0.1	0.3	-	0.2	0.4	0	2.6
30°N - 20°N		-	0.9	-	-	-	0.1	0	0.3	0.3	0.1	0.2	0	1.9
20°N - 10°N		-	-	-	-	0.2	0.8	-	0.2	1.3	2.8	-	-	5.3
10°N - 0°		0.6	-	0.4	-	0.8	1.8	0	0.1	1.7	1.7	0.2	0.2	7.5
0° - 10°S		2.3	-	-	-	0.9	2.7	0.1	0.1	0.2	1.0	0.1	0.3	7.7
10°S - 20°S		-	-	-	-	0	1.3	0.1	0.2	1.6	2.3	0.5	0.1	6.1
20°S - 30°S		-	-	-	-	-	0.3	-	2.0	0.9	1.3	0.3	0.2	5.0
30°S - 40°S		0.1	-	-	-	-	-	0.2	0.4	0.2	0.2	0.4	0.2	1.7
40°S - 50°S		-	-	-	-	-	-	-	0	-	-	-	-	0
50°S - 60°S		-	-	-	-	-	-	-	-	-	-	-	-	-
Total Emission		13.7	40.0	19.3	2.7	2.2	7.0	0.5	4.0	6.5	11.7	2.5	1.0	111.1